

# The concept of glomerular self-defense

MASANORI KITAMURA and LEON G. FINE

Department of Medicine, University College London Medical School, London, England, United Kingdom

**The concept of glomerular self-defense.** The balance between local offense factors and defense machinery determines the fate of tissue injury: progression or resolution. In glomerular research, the most interest has been on the offensive side, for example, the roles of leukocytes, platelets, complement, cytokines, eicosanoids, and oxygen radical intermediates. There has been little focus on the defensive side, which is responsible for the attenuation and resolution of disease. The aim of this review is to address possible mechanisms of local defense that may be exerted during glomerular injury. Cytokine inhibitors, proteinase inhibitors, complement regulatory proteins, anti-inflammatory cytokines, anti-inflammatory eicosanoids, antithrombotic molecules, and extracellular matrix proteins can participate in the extracellular and/or cell surface defense. Heat shock proteins, antioxidants, protein phosphatases, and cyclin kinase inhibitors may contribute to the intracellular defense. This article outlines how the glomerulus, when faced with injurious cells or exposed to pathogenic mediators, defends itself via the intrinsic machinery that is brought into play in resident glomerular cells.

In science, a new paradigm may arise from a tiny, unexpected finding that a researcher may confront in a routine experiment. The concept of “glomerular self-defense” occurred to us when we happened to meet an unpredicted experimental outcome.

In the initiation and progression of glomerulonephritis, it has been postulated that various cytokines play crucial roles. Once inflammation is initiated, glomeruli primed by infiltrating leukocytes may become more susceptible to injury by inflammatory cytokines. To examine this possibility, we tested the cytokine responses of normal and nephritic glomeruli using isolated glomeruli. As expected, when normal glomeruli were stimulated with a proinflammatory cytokine interleukin (IL)-1 $\beta$ , the expression of the cytokine-responsive metalloproteinase stromelysin was induced. Unexpectedly, however, the IL-1-triggered gene expression was blunted in glomeruli

in the regeneration phase of acute glomerulonephritis [1]. The blunted response was observed not only for stromelysin but also for other cytokine-inducible molecules. Furthermore, when activated macrophages were adoptively transferred into the nephritic glomeruli, the induction of inflammatory mediators in resident cells was suppressed, compared with the induction that occurred in normal glomeruli (abstracts; Ogura et al, *J Am Soc Nephrol* 8:487A, 1997; Sütö and Kitamura, *J Am Soc Nephrol* 8:481A, 1997) [1, 2]. We hypothesized that after an inflammatory insult, the glomerulus may acquire the potential for protecting itself from further activation and injury.

The idea of glomerular self-defense has not been addressed before, but the enhancement of local tissue defense in response to environmental stresses is a well-known phenomenon in certain pathophysiological situations. The best-known example is “ischemic preconditioning,” in which a brief period of ischemia results in the resistance of tissues to subsequent, severe ischemia. In the kidney, ischemic preconditioning prevents the reduction in glomerular filtration rate caused by ischemia/reperfusion or intra-arterial infusion of hydrogen peroxide. Another example of tissue defense is “thermotolerance.” In various tissues and cultured cells, the exposure to thermal stress induces a set of stress proteins, so-called heat shock proteins (HSPs), thereby affording tolerance against subsequent insults. It is therefore not surprising that the glomerulus, within inflammatory milieu, acquires an insensitivity to stimuli, although the acquisition of the “anti-inflammatory status” by inflamed organs or tissues is not well addressed to date. The aim of this review is to address possible mechanisms of local defense that may be exerted during glomerular injury. This article outlines how the glomerulus, when faced with injurious cells or exposed to pathogenic substances, defends itself via the intrinsic machinery that is brought into play in resident cells, especially in mesangial cells.

**Key words:** glomerulonephritis, cytokine inhibitor, anti-inflammatory cytokine, antioxidant, heat shock protein, cyclin kinase inhibitor.

Received for publication April 6, 1998


and in revised form July 20, 1998

Accepted for publication October 2, 1998

© 1999 by the International Society of Nephrology

## GENERAL PRINCIPLES AND CLASSIFICATION OF DEFENSE SYSTEMS

Many acute inflammations resolve without specific therapy. Like the initiation of inflammation, recovery from



**Fig. 1. Glomerular defense mechanisms: Defenders and their targets.** Abbreviations are: HSPG, heparan sulfate proteoglycan; SOD, superoxide dismutase.

inflammatory disease is a dynamic, multifaceted process in which a number of anti-inflammatory molecules are elaborated under well-ordered, switch-off programs. Figure 1 outlines glomerular defense against leukocytes, platelets/thrombi, and complement and inflammatory mediators. As illustrated, the vast array of factors can be mobilized in defense of the glomerulus, and its distribution and regulation is listed in Tables 1 and 2. A brief overview of the defense strategies is pertinent.

Toward resolution of inflammation, the defending machinery counteracts the attack by offenders at multiple levels, that is, extracellular, cell surface, and intracellular levels. For example, activated resident cells produce inactivators for leukocytes or platelets and inhibitors for specific mediators, including cytokines/growth factors, oxidants, proteinases, coagulants, and eicosanoids. A number of defending molecules are involved in this "extracellular defense." Complement-mediated injury is another important mechanism involved in the generation of glomerular disease. Glomerular cells have cell membrane-bound complement inhibitors that participate in the "cell-surface defense." In response to environmental

perturbation, resident cells produce intracellular stress proteins that reinforce cellular resistance to subsequent stresses. HSPs and antioxidant enzymes are involved in this "intracellular defense." In addition, several "self-cooling" devices are present in resident cells. These include the production of autocrine inactivators, the creation of new extracellular matrix (ECM), and the induction of protein phosphatases and cyclin kinase inhibitors (CKIs).

Endogenous defending molecules can be classified in different ways, based on the range of their targets and kinetics under pathophysiological situations. Some defending molecules have very specific targets. Thus, cytokine inhibitors, proteinase inhibitors, complement regulatory proteins, antithrombotic molecules, antioxidants, and CKIs are the typical examples of "man-to-man combat." On the other hand, other defending molecules have a wide range of targets. This may be viewed as "shotgun defense," which involves anti-inflammatory cytokines, anti-inflammatory eicosanoids, heparin/heparan sulfate proteoglycan (HSPG), and nitric oxide (NO). In pathological situations, the defense system is more complicated. Various defending molecules may form cascades

**Table 1.** Distribution of glomerular defenders

Defending molecule	Producer/Distribution					
	Mesangial cells	Epithelial cells	Endothelial cells	Leukocytes	GBM	Undetermined
<u>Extracellular/cell-surface defense</u>						
Cytokine inhibitors						
IL-1ra				○		▲
sTNFR				○		▲
SPARC	○▲	●				
decorin	○●					
Protein inhibitors						
TIMP	○					●
crystatin	○					
Complement inhibitors						
DAF	○▲					●
CR1		○●				
MCP	○●	○●	●			
CD59	○●	○●	○●			
Crry	○●	○	○●			
C1 inhibitor		○				
factor H	○●					
vitronectin	●		▲		●	
clusterin	○●	○●				
Nucleotidase						
ATP/ADPase					●	
Anti-inflammatory cytokines						
TGF-β1	○▲	○		○		●
IL-4	▲	▲		○▲		
IL-6	○●	○		○		
IL-10	○			○		▲
IL-13				○		▲
LIF	○			○		●
Anti-inflammatory eicosanoids						
prostaglandin E <sub>2</sub>	○●	○	○			
prostaglandin I <sub>2</sub>	○	○	○			
lipoxin A <sub>4</sub>	○					▲
lipoxin B <sub>4</sub>	○					
Anti-thrombotic molecules						
thrombomodulin	○●		●			
tissue factor pathway inhibitor	○					●
tissue-type PA	○	○	●			
urokinase-type PA		○●				
Extracellular matrix						
mesangial matrix	○●					
heparin/heparin sulfate proteoglycan	○●	○	○●		●	
Antioxidant						
extracellular-superoxide dismutase			●			
Other						
nitric oxide	○		●	○		
<u>Intracellular defense</u>						
Heat shock proteins						
HSP90		●				
HSP70	○	●				
HSP60						
HSP27	●					
heme oxygenase-1	○					
Antioxidants						
superoxide dismutases	○	○	○			●
catalase			○			●
glutathione peroxidase						●
Protein phosphatases						
MKP-1	○▲					
PAC1	○					
B23	○					
PTP <sub>γ</sub>		●				
RPTP-BK		●				
GLEPP1		●				
Cyclin kinase inhibitors						
p21		▲				
p27	○●	▲				

Symbols are: ○, expression/production in culture; ●, distribution in normal glomeruli; ▲, induction in diseased glomeruli.

Abbreviations are: GBM, glomerular basement membrane; IL-1ra, IL-1 receptor antagonist; sTNFR, soluble tumor necrosis factor receptor; SPARC, secreted protein acidic and rich in cysteine; TIMP, tissue inhibitor of metalloproteinases; DAF, decay accelerating factor; CR1, complement receptor 1; MCP, membrane cofactor protein; Crry, complement receptor related gene y; ATP/ADPase, adenosine triphosphatase and adenosine diphosphatase; TGF-β1, transforming growth factor-β1; LIF, leukemia inhibitory factor; PA, plasminogen activator; HSP, heat shock protein; MKP-1, MAP kinase phosphatase 1; PTP<sub>γ</sub>, protein-tyrosine phosphatase; RPTP-BK, receptor-type protein-tyrosine phosphatase in the brain and kidney; GLEPP1, glomerular epithelial protein 1; p21, p21<sup>Cip1/Waf1/Sdi1/Cap20</sup>; p27, p27<sup>Kip1</sup>.

**Table 2.** Regulation of glomerular defenders

Defending molecule	In vitro stimuli		In vivo situations	
	Up-regulation	Down-regulation	Up-regulation	Down-regulation
<u>Extracellular/cell-surface defense</u>				
Cytokine inhibitors				
IL-1ra	LPS, IL-4, TGF- $\beta$ (Mo/M $\Phi$ )		$\alpha$ -GBM	
sTNFR	LPS (Mo)		human GN	
SPARC	PDGF, EGF (Mes)		Heymann (Epi), $\alpha$ -Thy 1 (Mes)	
decorin	TGF- $\beta$ (Mes)		$\alpha$ -Thy 1 (Mes)	
Proteinase inhibitors				
TIMP	IL-1 $\beta$ , TNF- $\alpha$ , HG, TGF- $\beta$ (Mes)		human GN [IgA, DN], PAN	
Complement inhibitors				
DAF	complement (Mes)		human GN [IgA, MN, FGS, MPGN, HUS, SLE, crescentic] (Mes)	
CRI			human GN [MN, DN, IgA, SLE] (Epi)	
MCP	complement (Mes)		human GN [MPGN, SLE, IgA] (End, Mes)	
CD59	complement (Mes)		human GN [SLE] (End), human GN [MN]	
C1 inhibitor	IL-1 $\beta$ , TGF- $\beta$ (Epi)			
factor H	IFN $\gamma$ (Epi)			
vitronectin	IFN $\gamma$ (Mes)		human GN [IgA, HUS] (Mes), [MN] (End)	
clusterin	thrombin (Mes, Epi)		human GN [MN, etc.], Heymann	
Nucleotidase				
ATP/ADPase			endotoxemia, $\alpha$ -Thy 1, $\alpha$ -GBM	
Anti-inflammatory cytokines				
TGF- $\beta$ 1	Ang II, ANP, thrombin, TX		$\alpha$ -Thy 1, $\alpha$ -GBM, Habu, PAN, DN	
	HG, IC, stretch (Mes)		human GN [IgA, FGS, SLE, DN, crescentic]	
IL-4	HG, oxLDL (Epi)		$\alpha$ -GBM, human GN [IgA, SLE, MN, MCD]	
IL-6	LPS, IL-1 $\beta$ , TNF- $\alpha$ , GIP, Ang II, IC (Mes)		human GN [IgA, IgM, DN]	
	LPS, IL-1 $\beta$ , TNF- $\alpha$ (Epi)			
IL-10			$\alpha$ -GBM	
IL-13			$\alpha$ -GBM	
LIF			$\alpha$ -GBM	
Anti-inflammatory eicosanoids				
prostaglandin E <sub>2</sub>	LPS, IL-1 $\beta$ , TNF- $\alpha$ , PDGF (Mes)			
	IL-1 $\beta$ , TNF- $\alpha$ , IL-6, PDGF, Ang II, AVP		$\alpha$ -GBM, Heymann	
	ET-1, thrombin, ROI, NO, IC, HG (Mes)			
	AVP (Epi)			
prostaglandin I <sub>2</sub>	IL-1 $\beta$ , TNF- $\alpha$ (Mes)			
	AVP (Epi)			
lipoxin A <sub>4</sub>			$\alpha$ -GBM, ConA/ $\alpha$ -ConA	

or networks. That is, one defending molecule triggers the expression of another or synergistically cooperates with others against offenders. The “defense cascade” or “defense network” may be important for the defense programs of the glomerulus.

Many defending molecules are inducible in response to a local inflammatory milieu. This “inducible defense” contributes to the resolution of established inflammation. In contrast, some defending molecules are constitutively expressed and/or are present in the glomerulus. This “constitutive defense” is conceivably involved in the prevention of injury and the maintenance of integrity of the glomerulus. For example, complement regulatory proteins and heparin/HSPG may function as the constitutive defenders. In both inducible and constitutive examples, defending molecules are produced by resident glomerular cells. Alternatively, these molecules may be adoptively transferred from other sites via the circulation and may be stored within the glomerulus. This “adoptive defense” should also be considered in the pathophysiological contexts.

## SELF-DEFENSE MACHINERY IN THE GLOMERULUS

### Extracellular defense

#### Inhibitors of inflammatory mediators

During inflammation, a number of extracellular mediators are released from infiltrating cells and resident cells, leading to the magnification of inflammatory processes. However, in many cases, inflammation is self-limiting. This is due to intrinsic mechanisms designed to control inflammatory processes. Both resident and infiltrating cells have the ability to elaborate a set of natural inhibitors against specific, pathogenic mediators. This “man-to-man combat” system is widely observed in mammals and plays an important role in the prevention and resolution of inflammation. This section summarizes the potential roles of natural inhibitors that regulate the activity of pathogenic substances in the glomerulus.

#### (1) Inhibitors of cytokines and growth factors

(a) Interleukin-1 receptor antagonist (IL-1ra). Interleukin-1 is a proinflammatory cytokine that plays a crucial

Table 2. Continued

Defending molecule	In vitro stimuli		In vivo situations	
	Up-regulation	Down-regulation	Up-regulation	Down-regulation
Anti-thrombotic molecules				
thrombomodulin	<i>LPS, TNF-<math>\alpha</math> (isol.G)</i>		human GN [MPGN, SLE]	
tissue factor pathway inhibitor	heparin (Mes)		$\alpha$ -GBM	
tissue-type PA	IL-1 $\beta$ (Mes)		$\alpha$ -GBM, human GN [IgA, SLE]	
urokinase-type PA	IL-1 $\beta$ , TNF- $\alpha$ , thrombin (Epi)			
Extracellular matrix	TNF- $\alpha$ , thrombin (Epi)		<i>human GN [SLE]</i>	
mesangial matrix	IL-1 $\beta$ , TNF- $\alpha$ , TGF- $\beta$ , HG, TX		various human and experimental GN	
heparin/heparan sulfate proteoglycan	Ang II, AVP, oxLDL, stretch (Mes)			
Other	TGF- $\beta$ , <i>Ang II, HG</i> (Mes)		$\alpha$ -Thy 1, PAN, renal ablation	
nitric oxide	LPS, IL-1 $\beta$ , TNF- $\alpha$ , IFN $\gamma$ (Mes)		$\alpha$ -Thy 1, $\alpha$ -GBM, Heymann, <i>in situ</i> GN	
	<i>TGF-<math>\beta</math>, PDGF, MP, PGE<sub>2</sub> (Mes)</i>			
	bradykinin, ATP, thrombin, PAF (End)			
Intracellular defense				
Heat shock proteins				
HSP70	<i>IL-1<math>\beta</math> (Mes)</i>		PAN, $\alpha$ -GBM	
HSP27			PAN (Epi)	
heme oxygenase-1	IL-1 $\beta$ (Mes)			
Antioxidants				
superoxide dismutase	IL-1 $\beta$ , ROI, heat-aggregated IgG, MP (Mes)		injection of ROI, MP	
	IL-1 $\alpha$ , LPS, ROI (Epi)			
	ROI, MP (End)			
catalase	MP (End)		injection of ROI, MP	
glutathione peroxidase			injection of ROI, MP	
Protein phosphatases				
MKP-1	ROI (Mes)		$\alpha$ -Thy 1, $\alpha$ -GBM (Mes)	
GLEPP1			$\alpha$ -GBM (Epi)	
Cyclin kinase inhibitors				
p21			$\alpha$ -Thy 1 (Mes), Heymann (Epi)	
p27	TGF- $\beta$ , <i>PDGF, bFGF</i> (Mes)		$\alpha$ -Thy 1 (Mes), Heymann (Epi)	

Abbreviations are: LPS, lipopolysaccharide; Mo/M $\Phi$ , monocytes/macrophages; PDGF, platelet-derived growth factor; EGF, epidermal growth factor; Mes, mesangial cells; HG, high glucose concentration; Epi, glomerular epithelial cells; IFN $\gamma$ , interferon- $\gamma$ ; Ang II, angiotensin II; ANP, atrial natriuretic peptide; TX, thromboxane; oxLDL, oxidized low density lipoprotein; GIP, granulocyte inhibitory protein; IC, immune complex; AVP, arginine vasopressin; ET-1, endothelin-1; ROI, reactive oxygen intermediates; NO, nitric oxide; isol.G, isolated glomeruli; MP, methylprednisolone; PAF, platelet activating factor; End, glomerular endothelial cells; bFGF, basic fibroblast growth factor;  $\alpha$ -GBM, anti-glomerular basement membrane nephritis; GN, glomerulonephritis; Heymann, Heymann nephritis;  $\alpha$ -Thy 1, anti-Thy 1 glomerulonephritis; IgA, IgA nephropathy; DN, diabetic nephropathy; PAN, puromycin aminonucleoside nephrosis; MN, membranous nephropathy; FGS, focal/segmental glomerulosclerosis; SLE, lupus nephritis; MPGN, membranoproliferative glomerulonephritis; HUS, hemolytic uremic syndrome; crescentic, crescentic glomerulonephritis; Habu, Habu-venom-induced glomerulonephritis; MCD, minimal change disease; IgM, IgM nephropathy; ConA/ $\alpha$ -ConA, concanavalin A/anti-concanavalin A immune complex glomerulonephritis; *in situ* GN, *in situ* immune complex glomerulonephritis. Producers are in parentheses.

role in a wide range of inflammatory disorders. In glomerulonephritis, IL-1 has been regarded as one of pivotal pathogenic mediators. IL-1, released by inflammatory cells, causes resident cells to proliferate and stimulate aberrant matrix metabolism, expression of adhesion receptors, and the release of inflammatory mediators, including cytokines/growth factors, chemokines, bioactive lipids, metalloproteinases, and reactive oxygen/nitrogen species [3, 4]. *In vivo*, a number of studies have suggested a role for local IL-1 in the pathogenesis of experimental and human glomerulonephritis [4].

Interleukin-1 receptor antagonist is a naturally occurring IL-1 inhibitor originally identified as a protein secreted from myelomonocytic cells. This molecule is structurally homologous to IL-1 $\alpha$  and IL-1 $\beta$  and competes with IL-1 for binding to IL-1 receptors type I and II [5]. The therapeutic utility of IL-1ra has been reported in various experimental diseases [6]. In the kidney, systemic administration of IL-1ra suppresses: glomerular

cell proliferation, the expression of macrophage inflammatory protein-2 (MIP-2) and intercellular adhesion molecule-1 (ICAM-1), the formation of intracapillary thrombi, leukocyte infiltration (neutrophils, monocyte/macrophages, and/or lymphocytes), and proteinuria in anti-glomerular basement membrane (GBM) nephritis and anti-Thy 1 glomerulonephritis in rats [7–12]. A similar therapeutic effect was reported by the administration of soluble IL-1 receptor [10, 11]. Treatment with IL-1ra may be effective even after the establishment of chronic glomerulonephritis. Lan et al reported that the administration of IL-1ra halted the progression of established anti-GBM glomerulonephritis [13].

The expression of IL-1ra is induced in various inflammatory diseases. In normal rat glomeruli, the expression of IL-1ra is not detectable [14]; however, obvious up-regulation of IL-1ra is observed in glomeruli subjected to anti-GBM nephritis [14]. Currently, the cell type responsible for the expression is not determined. One pos-



sible candidate is inflammatory leukocytes [14] because neutrophils and monocyte/macrophages produce IL-1ra *in vitro*. The production of IL-1ra by resident glomerular cells should not be excluded because this molecule is also produced by nonleukocytic cells. Of note, the promoter region of the IL-1ra gene contains the nuclear factor- $\kappa$ B (NF- $\kappa$ B) binding site that may be activated in response to inflammatory stimuli [15]. Indeed, the expression of IL-1ra is up-regulated by IL-1, IL-4, IL-6, transforming growth factor- $\beta$  (TGF- $\beta$ ), and lipopolysaccharide (LPS) in monocyte/macrophages and hepatocytes [15–18]. Possibly, both resident and infiltrating cells produce IL-1ra, and locally produced IL-1ra may contribute to limiting the severity of glomerulonephritis.

(b) Soluble tumor necrosis factor (TNF) receptor (sTNFR). Similar to IL-1, TNF- $\alpha$  is a proinflammatory cytokine involved in the pathogenesis of glomerular diseases, including puromycin aminonucleoside (PAN) nephrosis, murine models of lupus nephritis, serum sickness nephritis, anti-GBM nephritis, and antineutrophil cytoplasmic autoantibodies-positive human glomerulonephritis [19]. TNF- $\alpha$ , which is mainly produced by infiltrating leukocytes, causes resident cells to express adhesion receptors for leukocytes and release of inflammatory mediators [19]. The blockade of the TNF action via neutralizing antibodies attenuates experimental glomerulonephritis [20, 21], suggesting its pathological contribution.

Soluble tumor necrosis factor receptors (sTNFRp55 and sTNFRp75) are naturally occurring TNF inhibitors, which are truncated forms of cell surface TNF- $\alpha$  receptors. These molecules, originally found in the plasma of clinical and experimental endotoxemia, neutralize TNF- $\alpha$ -induced cytotoxicity and immunoreactivity *in vitro*. The therapeutic utility of sTNFRs for glomerulonephritis has been reported by several investigators. For example, the systemic administration of sTNFR preserved renal function and suppressed histopathologic changes (intracapillary thrombi and crescent formation), expression of MIP-2 and macrophage migration inhibitory factor (MIF), neutrophil infiltration, and proteinuria in anti-GBM nephritis in rats [10, 11, 20, 22].

Urine of patients with chronic glomerulonephritis exhibits the TNF inhibitory activity. Suzuki et al identified that these inhibitors are sTNFRp55 and a homologue of sTNFRp75 [23]. This evidence raises possibilities that sTNFRs may be produced in nephritic glomeruli and that locally produced sTNFRs might contribute to limiting the action of TNF- $\alpha$  in the glomerulus. It has been shown that peripheral blood monocytes release sTNFR in response to LPS [24]. It has not been determined whether resident glomerular cells contribute to the production of sTNFR in nephritic glomeruli.

(c) Secreted protein acidic and rich in cysteine (SPARC). Platelet-derived growth factor (PDGF) is a prototypical

growth factor that plays an important role in various pathological situations. PDGF is expressed in a wide range of experimental and human glomerular diseases [25]. PDGF induces proliferation of mesangial cells *in vitro* and *in vivo* [25]. In anti-Thy 1 glomerulonephritis in rats, the inhibition of PDGF by the administration of an anti-PDGF neutralizing antibody resulted in the attenuation of cell proliferation and matrix expansion, suggesting the crucial role of PDGF in mesangial proliferative glomerulonephritis [26].

Secreted protein acidic and rich in cysteine, also known as osteonectin, is a glycoprotein produced by various cells. SPARC interacts with PDGF-BB and PDGF-AB and thereby inhibits their binding to PDGF receptors [27]. *In vitro*, the expression of SPARC is induced by growth factors including PDGF, TGF- $\beta$ , and insulin-like growth factor-I [28], suggesting possible up-regulation of this molecule under pathologic circumstances.

In the normal rat glomerulus, SPARC is constitutively expressed in glomerular epithelial cells [29]. This expression is increased in passive Heymann nephritis [29]. In contrast, after the induction of anti-Thy 1 glomerulonephritis, the expression of SPARC is induced in mesangial cells [30]. Cultured mesangial cells express and secrete SPARC, and the production is enhanced by PDGF and epidermal growth factor (EGF), the putative pathogenic mediators for anti-Thy 1 glomerulonephritis [30]. When SPARC is added externally, the DNA synthesis of mesangial cells is inhibited. Based on the role of PDGF in mesangial cell proliferation and matrix expansion *in vivo* [26], endogenous SPARC may be involved in self-defense of the glomerulus against inflammation.

(d) Decorin. Excessive deposition of ECM is the characteristic feature of tissue fibrosis. Numerous reports have suggested a role for TGF- $\beta$  in the fibrogenesis of organs. In glomerular disease, TGF- $\beta$  is up-regulated under various pathologic situations and has been regarded as a prosclerotic mediator [31]. TGF- $\beta$  induces the deposition of ECM by stimulating matrix production, decreasing ECM-degrading proteinases, and up-regulating proteinase inhibitors [32]. Although the role of TGF- $\beta$  in glomerular disease is not fully elucidated, these data suggest a possible contribution of TGF- $\beta$  to the expansion of ECM in nephritic glomeruli. Indeed, the administration of an anti-TGF- $\beta$  neutralizing antibody or an antisense TGF- $\beta$  oligodeoxynucleotide attenuated accumulation of ECM in acute anti-Thy 1 glomerulonephritis [33, 34].

Decorin is a proteoglycan that binds TGF- $\beta$ 1, TGF- $\beta$ 2, and TGF- $\beta$ 3 and, thereby, neutralizes their biological activities [35]. Border et al reported that systemic injection of recombinant decorin inhibited matrix expansion and proteinuria in acute anti-Thy 1 glomerulonephritis in which TGF- $\beta$  plays an important role [35]. Similarly, intramuscular transfer of a decorin gene effectively sup-

pressed the accumulation of glomerular ECM in the same experimental model [36].

Cultured mesangial cells constitutively produce decorin, and its production is up-regulated after exposure to TGF- $\beta$  [37]. Expression of decorin is induced in glomeruli in parallel with TGF- $\beta$  expression in anti-Thy 1 glomerulonephritis [38]. These data suggest a possibility that decorin is involved in an autoregulatory mechanism that limits excessive action of TGF- $\beta$  during acute glomerulonephritis.

## (2) Proteinase inhibitors

### (a) Tissue inhibitor of metalloproteinases (TIMP).

Matrix metalloproteinase (MMP) is a family of enzymes that degrades ECM components. The MMP family includes interstitial collagenases, gelatinases, stromelysins, and punctuated metalloproteinases (PUMPs) and plays an important role in the normal turnover of ECM. Aberrant activity of MMP is associated with various pathologies, including tissue fibrosis, inflammation, tumor invasion/metastasis, and atherosclerosis [39].

In the normal glomerulus, certain MMPs are constitutively expressed [40]. The glomerular expression of MMPs, including gelatinase A, gelatinase B, stromelysin-1, and PUMP-1, is increased under pathologic situations such as anti-Thy 1 glomerulonephritis, PAN nephrosis, passive Heymann nephritis, diabetic nephropathy, and several human glomerulonephritis (abstract; Miyazaki et al, *J Am Soc Nephrol* 6:902, 1995) [41–45]. The cells responsible for the production of MMPs are resident cells and inflammatory leukocytes [46]. Elaborated MMPs contribute to structural alteration of the mesangial matrix and GBM [40, 42, 47], leading to proteinuria [48].

Recent investigations disclosed a role of MMPs in the regulation of glomerular cell function, especially mesangial cell proliferation. The inhibition of constitutive expression of gelatinase A in mesangial cells led to depressed mitogenesis [49, 50]. Overexpression of stromelysin-1 in mesangial cells caused enhanced mitogenesis and migration within ECM [51]. These results suggest that certain MMPs function as autocrine triggers for mesangial cell activation. Potential mechanisms involved in the cellular activation by MMPs were reviewed previously [52].

From the viewpoint described above, the inhibition of excessive MMPs is beneficial for the attenuation of glomerular disease. The activity of MMPs is tightly regulated by their endogenous, specific inhibitors, TIMPs [39]. TIMP-1 and TIMP-2 are constitutively expressed and/or are secreted by cultured mesangial cells and isolated normal glomeruli [53, 54]. The expression of TIMP-1 in mesangial cells is up-regulated by inflammatory cytokines, including IL-1 $\beta$  and TNF- $\alpha$  [55], TGF- $\beta$  [56], or certain metabolic stimuli such as high glucose concentrations [57]. TIMP-1 is up-regulated in glomeruli of diabetic nephropathy and PAN nephrosis [42, 58]. TIMPs ex-

pressed in glomeruli may function as endogenous defenders that protect glomeruli from MMP-mediated injury.

(b) Cystatin. Cysteine proteinase is a diverse family of lysosomal enzymes with broad substrate specificity. These enzymes, including cathepsins B, H, and L, are generally located in lysosomes. However, a number of normal and tumor cells secrete these enzymes into the extracellular space. Although cysteine proteinases function optimally at acid pH values, they may have extracellular activity in acidic microenvironment. It has been shown that cathepsins B, H, and L are normally present in rat glomeruli [46]. These enzymes are also secreted by activated macrophages and neutrophils [59, 60], the major effector cells in glomerular disease. Endogenous cysteine proteinases present in glomeruli degrade intact GBM [61]. A role of cysteine proteinases in the pathogenesis of glomerular inflammation has been indicated by some investigators. The administration of specific inhibitors of cysteine proteinases (E-64, Ep475) reduced proteinuria in anti-GBM glomerulonephritis in rats. This therapeutic effect was correlated with a marked decrease in the local activity of cathepsins B and L [62, 63].

The activity of cysteine proteinases is inhibited by naturally occurring inhibitors, including cystatins [64]. The expression of cystatin C, the major member of the cystatin superfamily found in body fluid, is observed in various human tissues, including the kidney [65]. It has been reported that cystatin C is secreted by cultured mesangial cells [66]. Locally produced cystatin C could be involved in the protection of glomeruli from cysteine proteinase-mediated injury.

### (3) Complement regulatory proteins

Complement is a major mediator of tissue injury in various glomerulonephritis. Activation of complement occurs through the classical pathway and the alternative pathway; the former is triggered by antibody binding to antigens, and the latter is directly activated by microorganisms and damaged cells. The activation of complement attracts inflammatory cells and triggers resident cells to express adhesion receptors and to produce inflammatory mediators, including reactive oxygen intermediates (ROI), proteinases, and cytokines/chemokines, leading to glomerular injury [67]. The activation of complement is controlled by endogenous regulatory proteins present in plasma (C1 inhibitor, C4-binding protein, factor H, factor I, vitronectin, clusterin) and on the cell surface [decay accelerating factor (DAF), membrane cofactor protein (MCP), complement receptor 1 (CR1), membrane attack complex (MAC) inhibitory factor, complement receptor-related gene y (Crry)]. Complement regulatory proteins are normally present in the glomerulus, and under pathological circumstances, local levels of these proteins alter [68]. The functional inhibition of these molecules exacerbates pathological processes, indicating crucial roles for complement regulatory pro-

teins in the protection of glomeruli from antibody/complement-mediated injury.

(a) Decay accelerating factor. Decay accelerating factor inhibits complement activation by intervention in C3/C5 convertases via removal of C2b from C4b and Bb from C3b. DAF is a membrane-bound protein and is anchored to the plasma membrane by a carboxy terminal glycosyl-phosphatidylinositol (GPI). DAF is present in normal human glomeruli [69]. Modest expression of DAF mRNA may be detectable in normal glomerular cells [70]. In human glomerular disease, DAF is frequently detected in the mesangium, where C1q and C3 are colocalized [70, 71]. Using *in situ* hybridization, Abe et al reported that DAF mRNA is up-regulated in mesangial cells in IgA nephropathy and membranous nephropathy [70]. *In vitro*, mesangial cells have the ability to produce functional DAF, and the production and membrane expression are up-regulated by terminal complement components [72]. Antibody-mediated inhibition of DAF in glomerular cells enhances susceptibility to complement-induced cytotoxicity [69]. A recent study using *ex vivo* perfusion of human DAF-transgenic pig kidneys with human blood revealed that overexpression of DAF is sufficient to inhibit complement activation on the glomerular endothelium [73].

(b) Complement receptor 1. Similar to DAF, CR1 inactivates C3b and C4b. CR1 also functions as a cofactor for factor I, a plasma serine proteinase that cleaves C3b and C4b. In the glomerulus, CR1 is present exclusively on the surface of podocytes [68]. Consistent with this, glomerular epithelial cells in culture produce CR1 [74]. The expression of CR1 in podocytes is decreased in certain glomerular diseases, including membranous nephropathy, diabetic nephropathy, IgA nephropathy, and diffuse proliferative lupus nephritis [75–77]. The administration of a soluble form of CR1 lacking transmembrane and cytoplasmic domains significantly reduced both histopathologic and functional consequences of anti-Thy 1 glomerulonephritis, concanavalin A/anticoncanavalin A glomerulonephritis, and passive Heymann nephritis in rats [78].

(c) Membrane cofactor protein. Membrane cofactor protein (MCP) is a type 1 membrane glycoprotein that serves as a cofactor for factor I in inactivating C3b and C4b deposited on the cell membrane. This inactivation of C3b and C4b by MCP is associated with the protection of cells from autologous complement attack. In the normal human glomerulus, MCP is present on the surface of endothelial, mesangial, and epithelial cells [68]. Consistent with this, glomerular epithelial and mesangial cells in culture produce MCP [79]. The expression of MCP is up-regulated by activated complement in cultured mesangial cells [80]. MCP is increased in the glomerular capillary and mesangium of human kidney diseases, including membranoproliferative glomerulonephritis, IgA

nephropathy, and lupus nephritis [81]. Currently, the *in vivo* function of MCP is not fully determined.

(d) CD59. Like DAF, CD59 (also named MAC, protectin, or homologous restriction factor-20) is another GPI-anchored protein that inhibits the incorporation of C9 into C5b-9 complexes. In the normal glomerulus, CD59 is present on the surface of endothelial, mesangial, and epithelial cells [68]. Cultured glomerular epithelial, mesangial, and endothelial cells express CD59 [82, 83]. The expression of CD59 is up-regulated in glomerular epithelial cells by IL-1 $\beta$  and TGF- $\beta$  [84] and in mesangial cells by activated complement [85]. When treated with anti-CD59 antibody, these cells become more susceptible to complement-mediated lysis *in vitro* [82, 83]. In contrast, mesangial cells transfected with a CD59 cDNA show remarkable resistance to both lytic complement attack and sublytic effect of C5b-9 [86]. The expression of CD59 in the glomerular capillaries is increased in diffuse lupus nephritis [87]. In contrast, glomerular CD59 is decreased in membranous nephropathy [88].

*In vivo* roles of CD59 have been examined by several investigators. Matsuo et al examined the effect of anti-CD59 monoclonal antibody in a rat model of complement-dependent glomerulonephritis [89]. They found that local inhibition of endogenous CD59 enhanced glomerular cell proliferation, ICAM-1 expression, leukocyte influx, and fibrin deposition. Nangaku et al examined the effect of anti-CD59 antibody in the pathogenesis of thrombotic microangiopathy induced by an antibody to endothelial cells [90]. They found that local neutralization of CD59 exacerbated the disease, that is, more glomerular C5b-9 formation, endothelial damage, and platelet/fibrin deposition, resulting in the deterioration of renal function. These *in vivo* data clearly evidenced that CD59 is an endogenous defender against complement-mediated glomerular injury.

(e) Complement receptor-related gene y. In humans, DAF and MCP work at the step of C3/C5 convertases. Crry is the rodent analogue that possesses the function of DAF and MCP. In the normal rat glomerulus, Crry is present on the surface of mesangial and endothelial cells [68]. *In vitro*, all of the glomerular epithelial, mesangial, and endothelial cells produce Crry [82, 83]. When treated with an anti-Crry antibody, these cells become more susceptible to complement-mediated cytotoxicity [82, 83]. Furthermore, mesangial cells overexpressing Crry exhibit resistance to complement-mediated cell lysis and to sublytic stimulatory effects of complement [91].

*In vivo* roles of Crry have been investigated by a few investigators. Using a monoclonal antibody against Crry, Matsuo et al showed that Crry plays a vital role by preventing complement activation on the vascular endothelium [92]. Using the same strategy, they also showed that local inhibition of Crry resulted in accelerated mesangio-



lysis and leukocyte influx in anti-Thy 1 glomerulonephritis [93].

(f) C1 inhibitor. C1 inhibitor binds to C1 and C1r:C1s and releases them from C1q. It has been shown that glomerular epithelial cells in culture produce C1 inhibitor and that its production is enhanced by interferon- $\gamma$  [94]. The distribution of this molecule in the normal glomerulus remains to be determined.

(g) Factor H. Factor H binds to C3b and exhibits decay-accelerating activity and cofactor activity. The crucial role of this molecule in the maintenance of glomerular integrity was supported by evidence that inherited deficiency of factor H results in glomerulonephritis in pigs and humans [95, 96]. In normal human kidneys, factor H is localized in the mesangium [97]. Consistent with this, cultured mesangial cells produce factor H, and its production is enhanced by interferon- $\gamma$  [98].

(h) Vitronectin. Vitronectin, also called S-protein, is a component of ECM. This molecule binds to C5b-7 and inhibits the action of MAC. In the normal human glomerulus, vitronectin is present in the mesangium and GBM [99, 100]. Currently, it is not known whether the normal distribution of vitronectin is due to local production by glomerular cells or by the deposition of plasma protein. In glomerulonephritis, vitronectin is abundantly present in the immune deposit of the glomerular capillaries and mesangium and is colocalized with C3b and C5b-9 [99, 100].

(i) Clusterin. Human clusterin (SP-40,40) was first identified in glomerular immune deposits of a patient with membranous nephropathy [101]. Clusterin is part of the fluid phase MAC and functions as a complement inhibitor [102]. This molecule binds to C3, C8, and C9, inhibits generation of MAC, and thereby attenuates complement-mediated cytolysis. In a number of immune-mediated glomerulonephritis, clusterin is found in the glomerulus in conjunction with components of MAC [103–105].

Laping et al reported that clusterin mRNA and protein are detectable in normal rat glomeruli [106]. Clusterin mRNA is constitutively expressed in cultured podocytes and mesangial cells, and the expression is up-regulated by certain stimuli, including thrombin [106]. Using kidneys subjected to complement-dependent, passive Heymann nephritis, Saunders et al reported that the isolated kidneys perfused with clusterin-depleted plasma developed increased proteinuria, accelerated deposition of terminal complement components, and enhanced podocyte injury in the glomerulus [107]. This result indicates a possible role for local clusterin in the protection of glomeruli from complement-mediated injury.

#### (4) *Oxygen radical scavengers*

Extracellular superoxide dismutase (EC-SOD). Superoxide dismutases are mainly localized in the intracellular compartment but are also present in extracellular

fluid at low concentrations [108]. EC-SOD is a secretory glycoprotein that is the major SOD isozyme in extracellular fluids. The EC-SOD binds to heparan sulfate proteoglycan (HSPG) on endothelial cells and functions as a “protective coat” that attenuates oxidant-mediated cell injury [109].

#### (5) *Inhibitors of adenine nucleotides*

Adenosine triphosphatase and adenosine diphosphatase (ATP/ADPase). Extracellular adenine nucleotides, for example, ATP and ADP, have been considered as mediators of inflammation [110]. This is based on the fact that adenine nucleotides trigger platelet aggregation, production of ROI by neutrophils and macrophages, and proliferation of mesangial cells. The administration of adenine nucleotide analogues (ATP $\gamma$ S, ADP $\beta$ S) increases intraglomerular platelet aggregation and leukocyte production of superoxide in anti-Thy 1 glomerulonephritis [111]. In contrast, the administration of 2-chloro-adenosine inhibits intraglomerular platelet aggregation, leukocyte superoxide production, and proteinuria in the early phase of anti-Thy 1 glomerulonephritis [111]. These data indicate the proinflammatory activities of extracellular adenine nucleotides in the glomerulus. ATP and ADP are converted to adenosine by ATP/ADPase. ATP/ADPase is therefore potentially anti-inflammatory via the removal of proinflammatory ATP/ADP and via the generation of the anti-inflammatory molecule, adenosine.

In the glomerulus, ADPase activity is detected in the GBM [112]. The role of endogenous ADP/ATPase in glomerulonephritis has been extensively investigated by Bakker et al [110]. Depressed activity of glomerular ADPase is observed in experimental endotoxemia, anti-Thy 1 glomerulonephritis, and anti-GBM nephritis in which the procoagulant state is present [113, 114]. Poelsstra et al reported that impairment of ADPase alone did not induce platelet aggregation in the glomerulus, but facilitated it under certain prothrombotic conditions [115]. Using *ex vivo* perfusion of isolated kidneys with platelets and ADP, they also showed that selective inhibition of ADPase resulted in enhanced platelet aggregation in the glomerulus [116]. These data show that endogenous ADP/ATPase is a component of glomerular self-defense against platelet-mediated injury.

#### Inactivators of leukocytes and resident cells

One of the most common pathological features of glomerular disease is infiltration of leukocytes. These are mainly monocytes/macrophages, with neutrophils and T lymphocytes present in smaller numbers [117]. Several investigators have disclosed a link between leukocyte infiltration and structural/functional alteration of the glomerulus [118]. Using experimental glomerulonephritis in combination with leukocyte depletion, the roles of leukocytes in glomerular injury have been established.

During resolution of acute glomerulonephritis, acti-

vated leukocytes must be deactivated. Several mediators, including peptide factors and bioactive lipids, have been postulated as “deactivators” of inflammatory leukocytes. Some of these are produced in normal glomeruli, and the production is up-regulated under pathologic situations. At inflammatory sites, the cooperation of leukocyte inactivators may play a role in the prevention of or recovery from glomerulonephritis.

After the initiation of glomerulonephritis, leukocytes and platelets activate resident cells via releasing pathogenic mediators. Once activated, however, resident cells may elaborate a set of “self-inactivators” and thereby regulate their activity. This autoregulation may be important to control inflammatory processes in the glomerulus. The distribution and putative function of these anti-inflammatory mediators in the glomerulus are addressed below.

#### (1) Anti-inflammatory cytokines.

(a) Transforming growth factor- $\beta$ 1 (TGF- $\beta$ 1). Transforming growth factor- $\beta$ 1 is a pleiotropic, “Janus-faced” cytokine involved in a wide range of pathophysiological processes. Overexpression of TGF- $\beta$ 1 is closely linked to certain pathologies, including fibrogenesis [31]. On the other hand, there are lines of evidence that support the bright side of TGF- $\beta$ 1. TGF- $\beta$ 1 is known to be a potent regulator of immune systems and inflammatory processes, generally functioning as an endogenous anti-inflammatory molecule [119, 120]. TGF- $\beta$ 1 exerts anti-inflammatory actions in several diseases, especially in autoimmune disorders (abstract; Pankewycz et al, *J Am Soc Nephrol* 5:761, 1994) [121–124]. Constitutive expression of TGF- $\beta$ 1 is essential for the maintenance of normal immune and organ functions, as targeted disruption of TGF- $\beta$ 1 via homologous recombination induced immune dysregulation, multiorgan inflammation, and early death in mice [125–127].

Normal glomeruli express or contain detectable levels of TGF- $\beta$ 1 [38, 128–130]. The physiological meaning of this observation is currently unknown. Based on the fact that TGF- $\beta$  generally functions as a differentiation factor, the basal levels of TGF- $\beta$ 1 could contribute to the maintenance of normal glomerular structure and function. In various pathological conditions, TGF- $\beta$ 1 is up-regulated in the glomerulus. The induction of TGF- $\beta$ 1 has been reported in experimental models, including anti-Thy 1 glomerulonephritis, anti-GBM nephritis, Habu-venom glomerulonephritis, PAN nephrosis, and diabetic glomerulopathy, as well as in human diseases such as IgA nephropathy, focal/segmental glomerulosclerosis, crescentic glomerulonephritis, lupus nephritis, diabetic nephropathy, and human immunodeficiency virus nephropathy [31, 130]. The cell type responsible for TGF- $\beta$ 1 production in the glomerulus is not fully elucidated, but mesangial cells may be one of the major sources [38, 131]. Cultured glomerular epithelial cells

also produce TGF- $\beta$ 1 [132]. *In vitro*, mesangial cells have the ability to secrete latent TGF- $\beta$ 1 abundantly and to convert it to the mature form [133–135]. The production of TGF- $\beta$ 1 by mesangial cells and epithelial cells may be enhanced by a variety of factors, including angiotensin II, atrial natriuretic peptide, thrombin, thromboxane, high glucose, oxidized lipoprotein, immune complexes, and mechanical stress [132, 136–143]. In pathologic circumstances, TGF- $\beta$ 1, TGF- $\beta$ 2, and TGF- $\beta$ 3 may be similarly up-regulated in the glomerulus [130], but the pathophysiological role and kinetics of TGF- $\beta$ 2 and TGF- $\beta$ 3 remain to be disclosed.

In the generation of glomerulonephritis, macrophage-derived proinflammatory cytokines play crucial roles. Previous reports provided evidence that TGF- $\beta$ 1 inhibits production of IL-1 $\alpha$ , TNF- $\alpha$ , lymphotoxin, and interferon- $\gamma$  by peripheral blood mononuclear cells and peritoneal macrophages [144, 145]. We recently showed that cultured mesangial cells secrete a factor that strongly inhibits production of IL-1 $\beta$ , IL-6, TNF- $\alpha$ , and monocyte chemoattractant protein-1 (MCP-1) by activated macrophages. We have identified that this active entity is TGF- $\beta$ 1 [134, 146].

Reactive oxygen/nitrogen species play a role in a wide range of glomerular injury. In glomerulonephritis, major sources of ROI and NO are neutrophils and monocytes/macrophages [147, 148]. It is known that TGF- $\beta$ 1 inhibits production of ROI and NO by macrophages at picomolar concentrations [149, 150]. Locally produced TGF- $\beta$ 1, possibly, functions as a safeguard molecule that attenuates ROI/NO-mediated, glomerular injury.

TGF- $\beta$ 1 is an inhibitor of mitogenesis in cultured glomerular cells. Several investigators have shown that, *in vitro*, externally added TGF- $\beta$ 1 inhibits mitogenesis of cultured mesangial, epithelial, and endothelial cells [151–153]. Similarly, the proliferative response of isolated glomeruli is inhibited by externally added TGF- $\beta$ 1 [154]. Using an *ex vivo* gene transfer approach [155], we have demonstrated that mitogenic responses of glomerular cells in acute anti-Thy 1 glomerulonephritis could be repressed by the introduction of a gene coding for the active form of TGF- $\beta$ 1 [154].

In certain cell types, TGF- $\beta$  counteracts the effects of proinflammatory cytokines [156–158]. Previous reports showed that mesangial cells, endothelial cells, and isolated glomeruli treated with TGF- $\beta$ 1 exhibited depressed responses to IL-1 $\beta$  and TNF- $\alpha$  (abstract; Danoff and Jiang, *J Am Soc Nephrol* 7:1696, 1996) [154, 159, 160]. Mesangial cells stably transfected with a cDNA encoding the active form of TGF- $\beta$ 1 exhibit blunted expression of stromelysin in response to the macrophage cytokine IL-1 $\beta$  [1]. In contrast, mesangial cells transfected with a dominant-negative mutant of TGF- $\beta$  type II receptor show enhanced responses to IL-1 [1]. This result indicates that TGF- $\beta$  functions as an autocrine inhibitor of

the cytokine response in mesangial cells. In isolated rat glomeruli, externally added TGF- $\beta$ 1 repressed the induction of stromelysin by macrophage-derived cytokines. Similarly, when isolated nephritic glomeruli producing active TGF- $\beta$ 1 (anti-Thy 1 nephritis, day 7) were stimulated by IL-1 $\beta$ , the induction of stromelysin was depressed, compared with normal glomeruli [1].

We previously reported that *in vivo* transfer of activated macrophages induces the expression of stromelysin in resident glomerular cells [161]. To investigate whether endogenous TGF- $\beta$ 1 suppresses the macrophage-triggered glomerular cell activation, LPS-stimulated reporter macrophages were transferred into normal rat glomeruli, and glomeruli were subjected to anti-Thy 1 glomerulonephritis (day 7). Compared with normal glomeruli, the nephritic glomeruli secreted higher levels of biologically active TGF- $\beta$ . As described earlier in this article, in the normal glomeruli, stromelysin mRNA was markedly induced in resident cells after the transfer of activated macrophages. This induction was dramatically suppressed in the nephritic glomeruli producing active TGF- $\beta$ 1 [1]. Similar suppression was also observed in the induction of other cytokine-inducible molecules, including gelatinase B [2] and inducible NO synthase (iNOS) (abstract; Sütö and Kitamura, *ibid*). Based on this evidence together with the *in vitro* data, TGF- $\beta$ 1 can be regarded as a potential defender molecule that attenuates activation of resident cells faced by inflammatory macrophages.

**(b) Interleukin-4.** Interleukin (IL)-4 is a cytokine that is produced mainly by T-cell helper type 2 (Th2). It has been shown that IL-4-positive T cells are present in glomeruli of a rat model of anti-GBM nephritis and human glomerulonephritis, including IgA nephropathy and lupus nephritis [162, 163]. However, IL-4 may also be produced by resident glomerular cells. Using *in situ* hybridization, Furusu et al reported that IL-4 mRNA is induced in mesangial cells and podocytes in human glomerulonephritis, including IgA nephropathy, lupus nephritis, membranous nephropathy, and minimal change nephrotic syndrome [164].

Interleukin-4 is a potent inactivator of macrophages. IL-4 inhibits production of IL-1 $\beta$ , IL-8, and MIP-1 $\alpha$ , stimulates the expression of IL-1 receptor antagonist and IL-1 decoy receptor, and suppresses the production of ROI and NO in macrophages and neutrophils [165]. Tipping et al reported that the administration of IL-4 attenuated crescent formation, accumulation of T cells/macrophages and proteinuria in a murine model of anti-GBM nephritis [166]. Using IL-4 null mutant mice, Kitching et al recently showed that IL-4 deficiency amplified local and systemic Th1 responses and exacerbated anti-GBM glomerulonephritis [167]. Although there is some controversy [168, 169], locally produced IL-4 may participate in the attenuation of glomerular inflammation.

In addition to its effects on leukocytes, IL-4 modulates the function of resident glomerular cells. For example, IL-4 antagonizes the effect of IL-1 and inhibits proliferation of mesangial cells [170]. Cytokine induction of E-selectin in glomerular endothelial cells is also inhibited by IL-4 [171]. Endogenous IL-4 may attenuate inflammatory processes via inhibiting accumulation of leukocytes and cytokine responses of resident cells.

**(c) Interleukin-6.** Interleukin-6 is a multifunctional cytokine produced by various cell types and regulates growth and differentiation of a wide range of cells. Cultured glomerular epithelial and mesangial cells have the ability to produce IL-6, and the production is increased by inflammatory stimuli, including LPS, IL-1 $\beta$ , TNF- $\alpha$ , granulocyte inhibitory protein, angiotensin II, and immune complexes [172–177]. Expression of IL-6 may be detectable in the mesangium of normal human glomeruli [178]. The mesangial expression of IL-6 is markedly up-regulated during glomerular inflammation, especially mesangial proliferative glomerulonephritis [173, 178, 179].

Initially, IL-6 was considered to be a proinflammatory cytokine because IL-6 is an activator of lymphocytes and induces proliferation and MCP-1 expression by mesangial cells *in vitro* and *in vivo* [173, 180, 181]. Systemic administration with IL-6 exacerbated, but blockade of IL-6 receptor ameliorated murine lupus nephritis [182, 183], indicating a pathogenic role of IL-6 in certain types of glomerulonephritis. However, subsequent investigation pointed that this story is not simple. Some investigators reported that IL-6 did not induce or rather inhibited proliferation of mesangial cells *in vitro* and *in vivo* [184–187]. Rees et al reported that administration with IL-6 attenuated glomerular thrombosis, leukocyte infiltration/activation, expression of cytokines/chemokines, and proteinuria in nephrotoxic serum nephritis (abstract; Karkar et al, *J Am Soc Nephrol* 6:834, 1995) [11, 188]. The putative, anti-inflammatory properties of IL-6 are as follows: (a) up-regulation of circulating cortisol concentrations and (b) suppression of cytokine production (IL-1 $\beta$ , TNF- $\alpha$ , IL-8, MCP-1), induction of cytokine inhibitors (IL-1ra, IL-1 decoy receptor, soluble IL-1 receptor, sTNFR), and inhibition of ROI/NO production by macrophages [165]. Using IL-6 null mutant mice, a recent investigation clearly showed that IL-6 is an anti-inflammatory cytokine required for controlling local and systemic inflammatory responses [189]. In certain pathologic circumstances, endogenous IL-6 synthesized by resident cells may be involved in the glomerular self-defense against inflammation.

**(d) Interleukin-10.** Interleukin-10 is a multifunctional cytokine that is an important regulator of lymphoid and myeloid cell functions. IL-10 is produced mainly by Th2 cells and has profound effects on the morphology, phenotype, and production of bioactive substances by monocytes/macrophages. IL-10 inhibits macrophage produc-



tion of reactive oxygen/nitrogen species and cytokines, including IL-1 $\alpha$ , IL-1 $\beta$ , IL-8, TNF- $\alpha$ , granulocyte macrophage colony-stimulating factor (GM-CSF), and granulocyte colony stimulating factor (G-CSF) [190–192]. IL-10 enhances the synthesis of cytokine inhibitors, IL-1ra, and sTNFR [193, 194]. Several investigators reported the utility of IL-10 for the treatment of experimental glomerulonephritis. Tipping et al showed that, like IL-4, pretreatment with IL-10 attenuated crescent formation, accumulation of T cells/macrophages, proteinuria, and renal insufficiency in a murine model of anti-GBM nephritis [166]. Kitching et al showed that systemic administration of IL-10 in combination with IL-4 reduced the severity of anti-GBM nephritis in mice even after the establishment of the disease [195].

Interleukin-10 is produced in nephritic glomeruli [162]. The cells responsible for the production may be inflammatory leukocytes, but activated mesangial cells also produce IL-10 *in vitro* [196, 197]. Fouqueray et al reported a possibility that IL-10 functions as an autocrine inhibitor of cytokine production in mesangial cells [197]. Interestingly, they also observed that the production of IL-10 by mesangial cells was stimulated by the anti-inflammatory cytokine, TGF- $\beta$ . In contrast, Chadban et al recently reported that IL-10 may act as an autocrine stimulator of mesangial cell proliferation both *in vitro* and *in vivo* [196]. Further investigation will be required to determine the role for IL-10 in the regulation of glomerular cell activity.

(e) **Interleukin-13.** Interleukin-13 is an IL-4-like Th2 cytokine that modulates the function of monocytes/macrophages and B cells. IL-13 inhibits production of various proinflammatory cytokines (IL-1 $\alpha$ , IL-1 $\beta$ , TNF- $\alpha$ ), chemokines (IL-8, MIP-1 $\alpha$ ), and leukocyte growth factors (GM-CSF, G-CSF) by monocytes/macrophages [198]. IL-13 also inhibits production of NO by monocytes/macrophages and increases their expression of IL-1ra [198]. It has been shown that the expression of IL-13 mRNA was detectable in kidneys subjected to anti-GBM glomerulonephritis [199]. Currently, it is unclear whether resident glomerular cells produce IL-13.

(f) **Leukemia-inhibitory factor (LIF).** Leukemia-inhibitory factor is a pleiotropic cytokine produced by a wide range of cell types. Low levels of LIF expression are detectable in normal rat glomeruli [200]. Up-regulation of LIF is observed in nephritic glomeruli, including anti-GBM nephritis [200]. The role of LIF in the pathogenesis of glomerular inflammation is currently unclear, but Tang et al reported that systemic administration of recombinant LIF ameliorated experimental anti-GBM nephritis. That is, it reduced macrophage infiltration, local expression of cytokines, and proteinuria [200]. Cultured mesangial cells produce low levels of LIF. The transcriptional expression and protein synthesis are dramatically up-regulated in response to inflammatory stimuli, includ-

ing IL-1 $\beta$ , TNF- $\alpha$ , PDGF, and LPS [201]. Under certain pathological milieus, mesangial cell-derived LIF may function as a defending molecule against inflammation.

## (2) *Anti-inflammatory eicosanoids.*

Eicosanoids serve as important modulators for various pathophysiological processes in the glomerulus. Eicosanoids affect the function of leukocytes and resident cells and participate in the generation of glomerular disease, especially glomerulonephritis [202]. A majority of eicosanoids, including thromboxane A<sub>2</sub>, prostaglandin (PG) F<sub>2 $\alpha$</sub> , leukotrienes B<sub>4</sub>, C<sub>4</sub>, and D<sub>4</sub>, and 15-, 12- and 5-hydroxyeicosatetraenoic acids are regarded to be proinflammatory. These mediators trigger leukocyte infiltration, coagulation and proliferation and matrix production by resident cells [202]. On the other hand, some eicosanoids function as anti-inflammatory factors. These include prostacyclins (PGE<sub>2</sub>, PGI<sub>2</sub>) and lipoxins (LXA<sub>4</sub>, LXB<sub>4</sub>) [202, 203].

(a) **Prostacyclins (PGE<sub>2</sub>, PGI<sub>2</sub>).** Prostacyclins are powerful autacoids with potential effects on the immune system. Among these, PGF<sub>2 $\alpha$</sub>  acts as a proinflammatory mediator because it facilitates neutrophil chemotaxis [204]. In contrast, PGE<sub>2</sub> and PGI<sub>2</sub> are anti-inflammatory. PGE<sub>2</sub> suppresses functions of lymphocytes, natural killer cells, macrophages, and neutrophils [205–209]. Importantly, PGE<sub>2</sub> and PGI<sub>2</sub> inhibit neutrophil chemotaxis [210] and production of IL-1 $\beta$  and TNF- $\alpha$  by activated macrophages [209, 211]. In addition, PGI<sub>2</sub> inhibits platelet aggregation [212].

Prostacyclins modulate not only leukocyte/platelet function but also the activity of resident glomerular cells. For example, PGF<sub>2 $\alpha$</sub>  acts as a mitogen for cultured mesangial cells [213]. In contrast, PGI<sub>2</sub> inhibits mesangial cell proliferation [213], and PGE<sub>2</sub> represses production of type I and III collagens by mesangial cells [214].

It has been shown that cultured glomerular endothelial, epithelial, and mesangial cells have the ability to produce PGE<sub>2</sub> and PGI<sub>2</sub> [215–217]. Normal glomeruli produce PGE<sub>2</sub>, which is derived mainly from mesangial cells [218]. *In vitro*, the production of PGE<sub>2</sub> by mesangial cells is up-regulated by a variety of inflammatory mediators, including IL-1 $\beta$ , TNF- $\alpha$ , IL-6, PDGF, angiotensin II, arginine vasopressin (AVP), endothelin-1, thrombin, ROI, NO, and immunoglobulins [219–225]. The production of PGE<sub>2</sub> and PGI<sub>2</sub> by glomerular epithelial cells is increased by AVP [226]. *In vivo*, enhanced production of PGE<sub>2</sub> is observed in anti-GBM nephritis and passive Heymann nephritis [227, 228]. Based on the anti-inflammatory properties of PGE<sub>2</sub> and PGI<sub>2</sub>, these molecules may function as endogenous defenders against glomerular inflammation. Indeed, McLeish et al reported that systemic administration of PGE<sub>2</sub> resulted in attenuation of hypercellularity, immunoglobulin deposition, and proteinuria, as well as preservation of renal function in immune complex glomerulonephritis in mice [229, 230]. Poelstra et al reported that, in anti-Thy 1 glomerulone-



phritis, the PGI<sub>2</sub> analogue Iloprost inhibited intraglomerular platelet activation and reduced proteinuria [231].

(b) **Lipoxins (LXA<sub>4</sub>, LXB<sub>4</sub>).** Lipoxin is a family of eicosanoids with potential anti-inflammatory properties. Lipoxins counteract the pathogenic action of leukotrienes produced in various glomerular diseases [202, 203]. For example, LXA<sub>4</sub> and LXB<sub>4</sub> inhibit leukotriene B<sub>4</sub>-mediated neutrophil chemotaxis, adhesion to endothelial cells, and transmigration across the vascular wall [232, 233].

Increased production of LXA<sub>4</sub> is observed in certain glomerular diseases, including anti-GBM nephritis and concanavalin A-ferritin glomerulonephritis [234, 235]. Neutrophil-platelet interaction is considered to be the major pathway for the lipoxin formation [203]. That is, platelet lipoxygenase (12-LO) converts the neutrophil-derived leukotriene A<sub>4</sub> to lipoxins. However, Garrick et al reported that rat mesangial cells also have the ability to generate LXA<sub>4</sub> from leukotriene A<sub>4</sub> [236]. A subsequent study showed that 12- and 15-LO activity is present in mesangial cells and is responsible for the production of LXA<sub>4</sub> and LXB<sub>4</sub> [237]. Katoh et al reported that glomerular 12/15-LO mRNA is up-regulated in rat glomeruli with nephrotoxic serum nephritis [238]. Interestingly, this up-regulation was correlated with increased expression of IL-4, a well-known trigger for LO expression. *In vitro*, externally added LXA<sub>4</sub> and LXB<sub>4</sub> cooperatively inhibited adhesion of neutrophils to mesangial cells [239]. *Ex vivo* exposure of rat neutrophils to LXA<sub>4</sub> attenuated their recruitment to inflamed glomeruli following the induction of concanavalin A-ferritin glomerulonephritis [234]. Resident cell-derived lipoxins may play a regulatory role in limiting the accumulation of neutrophils during glomerular disease.

Leukotriene D<sub>4</sub> is potentially proinflammatory and mitogenic to cultured mesangial cells [240]. It has been shown that LXA<sub>4</sub> antagonizes the action of leukotriene D<sub>4</sub> on mesangial cells. This might be attributed to competition of LXA<sub>4</sub> with leukotriene D<sub>4</sub> for binding to cell surface [241].

### (3) *Apoptotic cells.*

Recently, Voll et al reported the possibility that apoptotic cells negatively regulate the function of monocytes/macrophages [242]. Monocytes incubated with apoptotic cells showed increased secretion of the anti-inflammatory cytokine IL-10 and decreased production of proinflammatory cytokines IL-1, IL-12, and TNF- $\alpha$ . This anti-inflammatory action of apoptotic cells was consistently observed, regardless of cell types or triggers for apoptosis. Fadok et al reported that phagocytosis of apoptotic neutrophils suppressed production of IL-1 $\beta$ , IL-8, GM-CSF, TNF- $\alpha$ , leukotriene C<sub>4</sub>, and thromboxane B<sub>2</sub> by activated macrophages [243]. In contrast, the production of anti-inflammatory mediators TGF- $\beta$ 1 and PGE<sub>2</sub> was up-regulated in the presence of apoptotic cells. In the

glomerulus, apoptosis of glomerular cells, especially mesangial cells, is observed in various glomerular diseases [244–248]. Apoptotic mesangial cells could play a role in subsidence of inflammatory macrophages in acute glomerulonephritis.

### (4) *Mesangial matrix.*

The mesangial matrix is a structural constituent that supports glomerular architecture. Recent investigation disclosed a possibility that the mesangial matrix may serve as an safety device that halts mesangial cell activation and dedifferentiation. Several lines of evidence supports this hypothesis. (a) Within three-dimensional ECM reconstituted *in vitro*, mesangial cells exhibit little mitogenic activity and lower levels of  $\alpha$ -smooth muscle actin ( $\alpha$ -SMA) expression compared with the cells on plastic [249–251]. (b) Cultured mesangial cells incorporated in “hillocks,” the three-dimensional structures of natural ECM created by mesangial cells, show depressed mitogenic activity and attenuated  $\alpha$ -SMA expression [251, 252]. (c) Following the destruction of the mesangial matrix by Habu venom, mesangial cells exhibit remarkable mitogenesis, which is gradually attenuated in parallel with the reconstruction of the mesangial matrix [253]. (d) Overexpression of matrix-degrading proteinases is observed in experimental and human glomerulonephritis, and it is associated with mesangial cell activation *in vivo* (abstract; Miyazaki et al, *ibid*) [41–45]. (e) Enhancement of matrix-degrading activity via transfection with a certain MMP gene or external supply with active MMP protein induces accelerated mitogenesis of cultured mesangial cells [49, 51]. (f) Inhibition of endogenous matrix-degrading activity via either antisense RNA, ribozyme, or a pharmacological inhibitor results in the suppression of mitogenic activity and  $\alpha$ -SMA expression in cultured mesangial cells [49, 50]. (g) Pharmacological inhibition of MMP *in vivo* attenuates mesangial cell proliferation in anti-Thy 1 glomerulonephritis [254]. These data suggest a role for the mesangial matrix in the suppression of mesangial cell activation.

The mechanisms involved in the regulatory effect of three-dimensional ECM on mesangial cell phenotype are not yet determined, but recent investigations have raised some possibilities. Marx et al found that three-dimensional cultures using a collagen gel resulted in a remarkable down-regulation of PDGF  $\beta$  receptors on the surface of mesangial cells [255]. We have reported that an ECM-initiated signal(s) suppresses activation of mesangial cells via inactivation of CARG box element, the *cis*-element required for mitogenesis and  $\alpha$ -SMA expression [251].

### **Inhibitors of platelet function and thrombogenesis**

Pathogenic roles of platelets in glomerular injury have been well documented. In addition to their contribution to thrombogenesis, platelet is an important source of various, potentially pathogenic mediators. These include

eicosanoids, growth factors, vasoactive amines, proteolytic enzymes, ROI, complement, adenine nucleotides, and so on [256]. The release of these molecules from activated platelets subsequently induces glomerular cell proliferation and ECM synthesis, facilitates leukocyte influx and immune complex deposition, and alters glomerular hemodynamics, leading to glomerular injury [256]. In this section, we address endogenous, glomerular machinery that attenuates platelet activation and thrombogenesis.

#### (1) *Thrombomodulin.*

Thrombomodulin is a glycoprotein found on endothelial cells. Thrombomodulin acts as a cofactor for the thrombin-catalyzed activation of protein C and inhibits thrombogenesis. Thrombin binding to thrombomodulin induces internalization of the complex with transport to lysosomes, leading to its degradation. This suggests that thrombomodulin functions as a "scavenger receptor" for thrombin. In the normal glomerulus, a small amount of thrombomodulin is present along the endothelium [257]. In membranoproliferative glomerulonephritis and lupus nephritis, the amount of endothelial thrombomodulin is increased [258]. In contrast, *in vitro*, thrombomodulin activity in normal glomeruli is repressed by inflammatory stimuli including TNF- $\alpha$  and LPS [259]. A recent report showed that, in addition to endothelial cells, mesangial cells have the ability to produce thrombomodulin both *in vitro* and *in vivo* [260]. Systemic administration of recombinant thrombomodulin inhibited glomerular fibrin deposition in experimental endotoxemia in rats [261], suggesting that locally produced thrombomodulin may serve as a defending molecule with anticoagulant activity.

#### (2) *Tissue factor pathway inhibitor (TFPI).*

Tissue factor pathway inhibitor is a factor Xa-dependent feedback inhibitor of the tissue factor-factor VIIa complex. Systemic administration of a two-domain TFPI analogue significantly reduced endotoxin-induced, massive fibrin deposition in the glomerulus [262]. Normally, TFPI is produced by glomerular cells [263]. Human mesangial cells synthesize TFPI *in vitro*, and its production is up-regulated by heparin [264]. Erlich et al provided evidence that TFPI is an endogenous defender that counteracts fibrinogenesis in certain glomerular disease [263]. They showed the following: (a) Glomerular TFPI synthesis was initially decreased in fibrin-dependent, crescentic glomerulonephritis. (b) *In vivo* inhibition of TFPI using an anti-TFPI antibody exacerbated glomerular fibrin deposition and renal impairment. (c) Infusion of recombinant human TFPI significantly reduced glomerular fibrin deposition, proteinuria, and renal impairment in this nephritis model.

#### (3) *Plasminogen activators.*

Thrombi, initially composed of aggregated platelets, are gradually replaced by fibrin. The fibrin is then di-

gested by fibrinolytic enzymes released from endothelial cells and other cell types. The fibrinolysis is mediated by conversion of a plasminogen to the active proteolytic enzyme, plasmin, via the tissue-type (t-PA) and urokinase-type (u-PA) plasminogen activators. Fibrin deposition is a characteristic feature of certain experimental and human glomerulonephritis. For example, in anti-GBM nephritis, fibrinolytic activity is reduced in parallel with fibrin deposition [265]. This is mainly due to decreased expression of t-PA and increased plasminogen activator inhibitor-I. It has been shown that the systemic administration of t-PA protected animals from glomerular fibrin deposition, crescent formation, and renal dysfunction in anti-GBM nephritis [266].

Isolated human glomeruli produce both t-PA and u-PA [267]. Immunohistochemical analysis showed that t-PA is present exclusively in the endothelium, but u-PA exists in the cytoplasm of glomerular epithelial cells [267]. The amount of t-PA is increased in certain human glomerular diseases, including IgA nephropathy and lupus nephritis [268]. In contrast, the amount of u-PA may be decreased in glomeruli of lupus-prone mice [269]. *In vitro*, mesangial cells produce t-PA, but not u-PA [270]. In contrast, glomerular epithelial cells in culture synthesize abundant u-PA and small amount of t-PA [271]. The production of these PAs is up-regulated by thrombin, IL-1, and TNF- $\alpha$  [271, 272].

Recently, the crucial role of local t-PA in the attenuation of glomerular inflammation was demonstrated using gene knockout mice [273]. In this study, crescentic glomerulonephritis was induced in mice in which the genes for PAs had been disrupted by homologous recombination. They found that the deficiency of t-PA exacerbated the glomerular injury. In contrast, the deficiency of u-PA did not significantly affect severity of the disease. These data imply that t-PA is the major PA that protects the glomerulus from crescentic glomerulonephritis.

### **Other factors**

#### (1) *Heparin and heparan sulfate proteoglycan (HSPG)*

Heparin is generally described as an anionic polysaccharide or a sulfated glycosaminoglycan with irregular sequences. Heparin has been considered a therapeutic agent for various glomerulonephritides. Systemic administration of heparin attenuates experimental and human glomerular diseases, including anti-GBM nephritis, Habu venom-induced injury, PAN nephrosis, remnant-kidney models, anti-Thy 1 glomerulonephritis, and lupus nephritis [274, 275]. The therapeutic effect of heparin is ascribed to its anti-inflammatory properties, that is, inhibition of coagulation/thrombosis, complement inactivation, suppression of leukocyte function, and inactivation of resident cells.

Antithrombin III inhibits the function of thrombin, factor Xa and factor IXa. This inhibition is markedly

enhanced by heparin and naturally occurring HSPG. HSPG is present on the luminal surface of endothelial cells and is the natural catalyst of the inhibitory effect of antithrombin III [276]. Heparin also inhibits platelet activation by neutrophils [277] and enhances endothelial production of NO, the potential suppressor of platelet aggregation and adhesion [278]. Additionally, heparin promotes fibrinolytic processes [279].

Heparin has the ability to inhibit complement activation *in vivo* [280, 281]. The inhibitory action is at multiple levels. Heparin inhibits C1 hemolytic function via interaction with and potentiation of C1 inhibitor [282]. C3 and C4 conversions by classical pathway activators are also inhibited by heparin [282]. Surface-associated heparin inhibits zymosan-induced activation of the alternative pathway by augmenting the action of complement regulatory proteins on particle-bound C3b [283].

Heparin has the potential to suppress the activity of pathogenic proteinases. For example, in vascular smooth muscle cells (SMCs), heparin inhibits phorbol ester-triggered induction of interstitial collagenase, gelatinase B, and stromelysin mRNAs, as well as the synthesis of these proteins [284]. We reported that, in mesangial cells, heparin selectively suppressed cytokine-mediated induction of stromelysin without affecting the expression of procollagen  $\alpha 1$  (IV), laminin B2, and TIMP [55]. Heparin also inhibits the activity of neutrophil-derived enzymes, including elastase and cathepsin G [285, 286], both of which are potentially pathogenic to the glomerulus [46].

Heparin and HSPG inhibit leukocyte rolling and adhesion on the endothelium [286, 287]. This is, in part, due to (i) the inhibitory effect of heparin on the function of L- and P-selectin, the endothelial adhesion receptors required for capture and rolling of neutrophils [288] and (ii) the down-regulation of the adhesion receptor CD11b on the surface of neutrophils [289]. *In vitro*, heparin inhibits both random migration and directed locomotion of neutrophils toward zymosan-activated serum [290]. *In vivo*, the intravenous administration of heparin diminishes the influx of neutrophils into the peritoneal cavity of thioglycollate-treated mice [288]. Heparin inhibits cellular aggregation, degranulation, and superoxide anion generation by activated neutrophils [291]. Heparin inhibits cytotoxicity exerted by neutrophils [292] and may induce their apoptosis [293]. Additionally, heparin also attenuates certain function of lymphocytes and macrophages [281, 294].

Heparin and HSPG have direct actions on resident glomerular cells, especially mesangial cells. Heparin and HSPG inhibit mitogenesis, migration, and expression of immediate early genes (*c-fos* and *egr-1*) in cultured mesangial cells [295–300] and increase their production of HSPG [301]. Recently, we found that heparin inhibits spontaneous apoptosis of podocytes in isolated rat glomeruli and oxidant-induced apoptosis in cultured mesan-

gial cells (abstract; Ishikawa and Kitamura, *J Am Soc Nephrol* in press) [302]. Heparin and HSPG may also alter the function of glomerular epithelial and endothelial cells. These molecules inhibit proliferation of glomerular epithelial cells and suppress their responses to EGF [303, 304]. Heparin facilitates release of EC-SOD from endothelial cells, leading to their protection from ROI-mediated injury [305, 306].

Heparin-like species are widely distributed in the glomerulus, especially in GBM and the mesangial matrix [307, 308]. The amount of HSPG in the glomerulus is increased in experimental glomerular diseases, including anti-Thy 1 glomerulonephritis, PAN nephrosis, and renal ablation models [309]. *In vitro*, glomerular cells produce heparin-like glycosaminoglycans. For example, glomerular epithelial and endothelial cells secrete heparin-like molecules, which suppress mesangial cell proliferation [295, 296]. Mesangial cells and epithelial cells also produce HSPG, which functions as an autocrine or paracrine growth inhibitor [297, 304, 310, 311]. Interestingly, angiotensin II, a well-recognized pathogenic mediator for glomerular disease, suppresses production of HSPG, but TGF- $\beta 1$ , a putative defending molecule against acute glomerular injury, stimulates the production of HSPG in mesangial cells [312, 313]. The intercellular network via heparin-like molecules may be an important defense mechanism that protects resident cells from activation and injury.

## (2) Nitric oxide (NO).

The L-arginine-NO pathway is involved in the physiological function of mammalian organs, including the kidney. In the glomerulus, NO synthase is constitutively expressed, and the basal generation of NO participates in the regulation of glomerular hemodynamics [314].

Locally produced NO may function as an oxygen radical scavenger in the glomerulus. In the presence of NO, superoxide produced by activated cells is rapidly combined with NO. This reaction results in superoxide scavenging, leading to the mitigation of its cytotoxic effect [315]. Glomerular cells, especially mesangial cells, have the ability to express iNOS and synthesize NO in response to proinflammatory stimuli, including IL-1 $\beta$  and TNF- $\alpha$  [316, 317]. Sandau, Pfeilschifter and Brune reported that exogenously supplied superoxide induced apoptosis of mesangial cells and that NO inhibited the cytotoxic effect of superoxide anion [318].

Nitric oxide is an endogenous inhibitor of leukocyte function, acting particularly on neutrophils. Kubes, Suzuki and Granger reported that suppression of endothelial production of NO facilitated adhesion of neutrophils to the endothelium via up-regulation of CD11/CD18 expression [319]. NO also functions as an inhibitor of neutrophil superoxide production via direct action on the NADP oxidase [320]. It has been shown that NO released



by resident cells protects the cells from neutrophil-induced cytotoxicity [321]. These results suggest that endothelium-derived NO is an important endogenous inhibitor of leukocyte adherence and cytotoxic action.

Nitric oxide inhibits platelet adhesion to the damaged endothelium and platelet aggregation [322, 323]. Shultz and Raij showed that endogenous NO prevents endotoxin-induced glomerular thrombosis in rats [324]. This was further supported by Westberg et al, who showed that the administration of NO inhibitor deteriorated, but NO donor prevented glomerular thrombosis in a rodent model of endotoxemia [325].

Overproduction of NO in the glomerulus is supposed to be proinflammatory via induction of glomerular cell injury [326]. However, subtoxic levels of NO exert some beneficial actions on resident cells. For example, externally supplied NO suppresses proliferation, ECM production, and expression of ICAM-1 in cultured mesangial cells [139, 327–329]. NO also suppresses oxidant-initiated apoptosis of mesangial cells via scavenging superoxide anion [318]. NO may inhibit leukocyte–endothelial interaction via down-regulation of adhesion receptors on the surface of endothelial cells [330]. Recently, Haberstroh et al showed that endogenous NO suppresses LPS-triggered expression of Rantes in the glomerulus [331].

*In vitro*, the L-arginine-NO pathway is constitutively active in glomerular endothelial cells. The activity may be up-regulated by inflammatory mediators, including bradykinin, extracellular 5' adenosine triphosphate (ATP), thrombin, and platelet-activating factor [148]. In cultured mesangial cells, the L-arginine-NO pathway is inducible in response to proinflammatory stimuli, including LPS, IL-1, TNF- $\alpha$ , and interferon- $\gamma$  [148]. In certain pathologic circumstances, endogenous NO may function as a part of the self-defense program in the glomerulus.

## Intracellular defense

### Heat shock proteins

Heat shock protein is a family of highly conserved proteins synthesized under various pathophysiological circumstances. These molecules are classified into five major subfamilies: 110 kDa HSP (HSP110), 90 kDa HSP (HSP90), 70 kDa HSP (HSP70), 60 kDa HSP (HSP60), and low-molecular weight HSP [332]. Generally, constitutively expressed HSPs control maturation and turnover of intracellular proteins and thereby play a role in the maintenance of cellular integrity. On the other hand, inducible HSPs are synthesized in response to environmental perturbations, including ischemia, inflammation, oxidative stress, and toxin exposure and function as defending molecules that protect tissues from further injuries [332]. Recent investigations have disclosed that several HSPs are expressed in normal kidneys and that their expression pattern and levels are altered after pathologic insults [333].

#### (1) HSP70 family.

The HSP70 family includes two major members: a 73 kDa protein that is constitutively expressed and a 72 kDa protein that is inducible in response to stresses [332]. Constitutive HSP73 participates in intracellular protein targeting, processing and transport, and thereby plays an essential role in the maintenance of cell function. In response to a wide range of stress, HSP72 is rapidly synthesized and acts as a molecular chaperone; that is, with HSP60, this molecule binds to denatured or unfolded proteins and subsequently renatures or refolds the damaged proteins [334, 335]. This allows the cells to be protected against further injury and facilitates their recovery from established damage.

In normal rat glomeruli, HSP70 is expressed in podocytes and Bowman's epithelium [336]. Glomerular expression of HSP70 is up-regulated in mesangial cells in PAN nephrosis [336]. Currently, the expression and kinetics of HSP70 in human glomerular diseases remain to be elucidated. *In vitro*, cultured mesangial cells constitutively express HSP70 [337]. Mesangial cells transfected with a HSP70 cDNA acquire resistance to oxidant-induced apoptosis [337]. This protective effect might be due, in part, to the ability of HSP70 to suppress activation of c-Jun N-terminal kinase (JNK) [338], a putative mediator for oxidant-induced apoptosis in mesangial cells [339].

In certain pathological circumstances, the constitutive expression of HSP70 in the glomerulus is down-regulated. We found that glomerular expression of HSP70 was suppressed in rat kidneys subjected to anti-GBM glomerulonephritis. The down-regulation of HSP70 was caused, at least in part, by IL-1 and was associated with the incidence of glomerular cell apoptosis *in vivo* [337]. Similarly, mesangial cells treated with IL-1 $\beta$  showed depression of HSP70 and an increase in susceptibility to oxidant-triggered apoptosis. Transfection of mesangial cells with a HSP70 cDNA overcame the apoptosis-priming effect of IL-1 $\beta$  [337]. Depression of HSP70 by inflammatory cytokines may magnify glomerular injury.

#### (2) HSP60 and HSP90 families.

The HSP60 family of molecules acts as a molecular chaperone that facilitates proper maturation of newly synthesized proteins [334, 335]. The members of this family facilitate folding of monomeric proteins and catalyze the higher ordered assembly of oligomeric complexes [332]. HSP60 is constitutively expressed in mammalian kidneys [340], but its intrarenal distribution is not determined.

HSP90 is an abundant cytosolic protein, and its expression is up-regulated by certain stresses. This molecule is now regarded as a chaperone involved in signal transduction. HSP90 interacts with a large number of other cellular proteins, including tyrosine kinases and steroid receptors and thereby regulates the kinetics and activity of these signaling proteins [335]. In the normal glomerulus,



HSP90 is present in podocytes [341]. This constitutive expression may be involved in physiological functions of the cells via interacting with particular target proteins.

Currently, the contribution of HSP60 and HSP90 to glomerular self-defense is unknown. Recent reports suggested the possibility that these molecules could be involved not in the prevention, but in the generation of diseases. For example, HSP60 may serve as an antigenic determinant to which cytotoxic T cells respond [342]. In certain cell types, increased HSP90 may rather enhance cellular damage [343]. Further investigation is needed to disclose the distribution, kinetics, and pathophysiological roles of these HSPs in the glomerulus.

### (3) HSP27.

Like HSP70, HSP27 participates in cellular defense. Overexpression of HSP27 leads to reinforcement of cellular resistance to stresses, including heat shock and ROI [332]. In human kidneys, the expression of HSP27 is observed predominantly in collecting ducts and modestly in proximal/distal tubules and mesangial cells [344].

In the rat, HSP27 is localized in glomerular epithelial cells [345]. HSP27 may be involved in physiological function of podocytes. In PAN nephrosis, the effacement of foot processes is closely associated with the induction and phosphorylation of HSP27 [345]. The structure of foot process—an essential part of the filtration barrier—is controlled by actin microfilaments, and actin polymerization is known to be regulated by HSP27. The altered activity of HSP27 in podocytes might therefore cause aberrant foot process structure, leading to the development of proteinuria.

### (4) HSP32 (*heme oxygenase-1*).

Heme oxygenases are microsomal enzymes that catalyze the rate-limiting step in the degradation of heme to biliverdin, free iron, and carbon monoxide. Two isoforms of heme oxygenase have been identified. Heme oxygenase-1 is the stress-inducible form that is also termed HSP32, and heme oxygenase-2 is the constitutive isoform that is present under physiological conditions [346]. Various environmental factors trigger expression of heme oxygenase-1. These include heat shock, heme compounds, heavy metal ions, cytokines, and oxidative stress [346]. In particular, the induction of heme oxygenase-1 is regarded as a general response to oxidant stress in mammalian cells [347] and is implicated in a defense mechanism against ROI [348]. Indeed, transfection with a heme oxygenase-1 cDNA confers resistance of epithelial cells to oxidant stress [349].

Heme oxygenase-1 is induced under inflammatory situations. The role of heme oxygenase-1 in the inflammatory milieu is not fully determined, but recent investigations showed that heme oxygenase-1 attenuates inflammatory responses [350, 351]. Expression of heme oxygenase-1 in the glomerulus has not been reported, but *in vitro*, cultured mesangial cells express this enzyme abundantly

in response to IL-1 $\beta$  [352]. Together with the fact that cytokines and reactive oxygen/nitrogen species are important pathogenic mediators in glomerulonephritis [147, 314, 353], heme oxygenase-1 may be induced in inflamed glomeruli and could participate in the modulation of inflammatory processes.

## Antioxidant enzymes

Oxidant stress is involved in various diseases, including inflammation. It is supposed that ROI play important roles in the generation of a broad array of human and experimental glomerular diseases [147].

ROI released by local neutrophils and macrophages trigger glomerular cell death [337], degradation of ECM [47], formation of thrombi and, thereby, contribute to the generation of diseases [147]. ROI may induce massive proteinuria via causing glomerular sieving defect without apparent histologic abnormality [354]. ROI also participate in the activation of resident cells. This is due to the fact that a variety of signaling molecules are redox-sensitive [355, 356]. For example, in mesangial cells, ROI trigger calcium signaling and phosphorylation of tyrosine kinases and activate transcription factors including activator protein 1 (AP-1) and NF- $\kappa$ B [339, 357, 358]. ROI may therefore affect a diverse range of cell function, including proliferation, apoptosis, matrix metabolism, and production of inflammatory mediators [359, 360]. In particular, ROI stimulate the production of chemokines and the expression of adhesion receptors by resident cells [361, 362] and, thereby, attract circulating leukocytes at affected sites. Based on these data, endogenous antioxidant enzymes should be important as the self-defense machinery that prevents glomerular injury [363].

### (1) *Superoxide dismutases*.

Mn-SOD and Cu/Zn-SOD are included in this category. These SODs promote dismutation of the superoxide anion, an immediate oxygen metabolite with a high biological activity, to yield hydrogen peroxide. Mn-SOD is inducible by a variety of stimuli, including ROI, endotoxin, and inflammatory cytokines. In contrast, the expression of Cu/Zn-SOD is constitutive [363, 364]. Studies using transgenic mice provided evidence that overexpression of either SOD confers resistance to oxidant-induced injury [365, 366].

Mn-SOD is expressed in isolated normal glomeruli and in cultured mesangial, endothelial, and epithelial cells [367–369]. The role of glomerular SOD has been extensively investigated in PAN nephrosis, an experimental model of minimal change disease. In this model, the administration of SOD effectively attenuated proteinuria and foot process fusion of podocytes [370, 371]. Experimental depletion of SOD by the administration of diethylthiocarbamate aggravated the disease (abstract; Hara et al, *Proc XIth Int Congr Nephrol*, 1990, p 442),

indicating the protective role of endogenous SOD in PAN nephrosis.

The expression of Mn-SOD in cultured glomerular cells is up-regulated by hydrogen peroxide, heat-aggregated IgG, or IL-1 [369, 372, 373]. This implies that Mn-SOD is up-regulated during glomerulonephritis. Currently, however, information is very limited regarding the kinetics and roles of SOD in glomerular inflammation. One report showed that when SOD was administered intravenously in rats, nephrotoxic serum nephritis was attenuated [374].

#### (2) *Catalase and glutathione peroxidase.*

Catalase and glutathione peroxidase catalyze the decomposition of hydrogen peroxide to yield oxygen and water [147]. When overexpressed, either enzyme confers cellular resistance to oxidant-induced injury [375, 376].

Catalase and glutathione peroxidase are normally expressed in the glomerulus, and the expression is up-regulated by oxidant stress [367]. The administration of catalase is effective for the treatment of PAN nephrosis [371]. Experimental depletion of endogenous glutathione peroxidase by a selenium-deficient diet or administration of diethyl maleate aggravates PAN nephrosis (abstract; Miyai et al. *Proc XIth Int Congr Nephrol*, 1990, p 1442) [377]. These data suggest that endogenous catalase and glutathione peroxidase function as defending enzymes against oxidant-mediated glomerular injury. Of note, rats fed with a diet deficient in selenium and vitamin E develop glomerular disease characterized by proteinuria and reduced glomerular filtration rate [378]. This result implies a "housekeeping" role of glutathione peroxidase (and other antioxidant enzymes) in the maintenance of normal glomerular function.

#### (3) *Heme oxygenases.*

Heme oxygenase-1 and heme oxygenase-2 operate as antioxidants at two different stages. These enzymes decrease the levels of the potential pro-oxidants heme and heme proteins and increase tissue concentrations of the antioxidant, bile pigments [348]. The potential role of these enzymes in glomerular disease has been addressed in the section about heat shock proteins.

### **Protein phosphatases**

Phosphorylation of signaling proteins plays a cardinal role in the regulation of a number of cellular processes. The reversible control over protein phosphorylation requires not only protein kinases but also protein phosphatases. In many cases, target proteins are phosphorylated at specific sites by one or more protein kinases, leading to their activation. Following the phosphorylation, the phosphates are subsequently removed by specific protein phosphatases, resulting in deactivation of the target proteins. This allows for reversible and tight control over the activity of signaling proteins and halts cellular activation

[379]. In this regard, protein phosphatases may serve as a piece of intracellular defense mechanisms.

A number of protein phosphatases have been identified. These include protein-tyrosine phosphatases (PTPs), protein-serine/threonine phosphatases, and dual-specificity phosphatases. Currently, distribution, kinetics, and function of protein phosphatases in the glomerulus are largely unknown, but recent investigation has shed light on putative roles of some protein phosphatases. The family of mitogen-activated protein kinases (MAP kinases), including extracellular signal-regulated kinase (ERK), JNK, and p38 MAP kinase, plays a crucial part in cellular activation in response to a broad array of extracellular stimuli [380]. Dual-specificity protein phosphatases participate in dephosphorylation of MAP kinases and thereby attenuate cell proliferation and stress-induced gene expression [379]. In cultured mesangial cells, several dual-specificity phosphatases are constitutively expressed, that include MAP kinase phosphatase 1 (MKP-1), PAC1, and B23 [381]. Treatment of mesangial cells with vanadate, an inhibitor of dual-specificity phosphatases, triggers mitogenesis and growth factor production, suggesting a role of protein phosphatases in mesangial cell inactivation [382]. The expression of MKP-1 in cultured mesangial cells is up-regulated by ROI [383]. In the oxidant-mediated anti-GBM glomerulonephritis, expression of MKP-1 is rapidly induced in mesangial cells [383]. Under pathological situations, the up-regulated MKP-1 may function as a breaking mechanism that attenuates activation of glomerular cells.

Interestingly, several PTPs, including PTP $\lambda$ , receptor-type PTP in the brain and kidney (RPTP-BK) and glomerular epithelial protein 1 (GLEPP1), are constitutively and predominantly expressed in podocytes [384–386]. These PTPs may function as "housekeeping brakes" for podocytes and might explain their resistance to mitogenic stimuli.

### **Cyclin kinase inhibitors**

Proliferation of eukaryotic cells is positively controlled by a set of nuclear proteins, called cyclins and cyclin-dependent kinases (CDKs). Cyclins are periodically expressed during specific phases of the cell cycle, and bind to and activate specific CDKs. G1/S phase cyclins and CDKs are required for G1/S transition and DNA synthesis. G2/M phase cyclins are required for mitosis [387]. The activity of CDKs is negatively regulated by CKIs that bind to and inactivate specific CDK-cyclin complexes and thereby arrest the cell cycle [388]. There are two families of CKI: the Cip/Kip family, and the INK4 (inhibitor of cdk4) family. The former comprises p21<sup>Cip1/Waf1/Sdi1/Cap20</sup> (p21), p27<sup>Kip1</sup> (p27) and p57<sup>Kip2</sup>, and the latter does p15, p16, p18, and p19 [387]. The Cip/Kip family members inhibit function of cyclin-CDK complexes in the G1 and S phases. The INK4 family members inhibit cyclin-CDK

complexes only in the G1 phase. In various cell types, the overexpression of P21, p27, or P57 is sufficient to induce cell cycle arrest in G1.

Quiescent mesangial cells constitutively express p27 *in vitro*. Shankland et al investigated the role of p27 in proliferation of cultured mesangial cells [389]. After stimulation with PDGF or basic fibroblast growth factor, mesangial cells exhibited a substantial decrease in the p27 levels. Consequently, there was a marked increase in expression of cyclin A and CDK2. The decline of p27 was prevented by TGF- $\beta$ 1 that suppressed mesangial cell proliferation. Furthermore, antisense oligonucleotides against p27 enhanced growth factor-induced cell proliferation, suggesting the importance of p27 as an endogenous inhibitor of mesangial cell growth *in vitro*.

Shankland et al further examined the expression of cell-cycle proteins *in vivo* in normal rat kidneys and kidneys subjected to anti-Thy 1 glomerulonephritis [390]. They found that, in normal glomeruli, p27 was highly expressed, whereas the expression of p21 was low. In the nephritic glomeruli, the induction of mesangial cell proliferation was associated with increased expression of cyclin A and CDK2 and transient reduction in p27 levels. During the recovery phase, the expression of p21 was up-regulated and maintained until at least day 10, whereas the depressed levels of p27 returned to the baseline levels within five days. This result suggests that p27 and p21 play distinctive roles; that is, the former maintains mesangial cells to be quiescent, and the latter inhibits its ongoing proliferation.

The role of p27 in the glomerulus has been further investigated using p27-null mutant mice [391]. In the normal condition, the null mutant mice did not exhibit any abnormality in the glomerulus. However, when anti-GBM nephritis was induced, the mutant mice developed more severe disease—increased cell proliferation, apoptosis, and matrix accumulation—compared with wild-type mice [391]. This result suggested that p27 functions as a “safeguard molecule” of the glomerulus against inflammation.

In nephritic glomeruli, podocytes are relatively resistant to mitogenic stimuli compared with mesangial and endothelial cells. To determine the mechanisms responsible for the lack of podocyte proliferation *in vivo*, Shankland et al examined the expression of CKIs in passive Heymann nephritis, in which the podocyte is the target of complement-mediated injury [392]. Following antibody deposition and complement activation, marked up-regulation of p21 and p27 and subsequent inhibition of CDK2 activity were observed in podocytes.

Taken together, p21 and p27 function as endogenous brakes to inhibit or terminate proliferation of resident cells in response to pathogenic, growth stimuli.

## STRATEGIES FOR REINFORCEMENT OF GLOMERULAR SELF-DEFENSE

Pharmacological agents that attenuate functions of platelets/coagulants, leukocytes, complement, and hemodynamic stress have been considered for therapeutic intervention in glomerular disease. As an alternative strategy, the reinforcement of “self-defense” may be useful for attenuation of glomerular injury. Several tactics are available for this purpose. These include preconditioning, pharmacological modulation, and genetic manipulations.

### Preconditioning

A brief period of ischemia leads to resistance of various organs to subsequent, severe ischemia. Yoshioka et al reported that in the kidney ischemic preconditioning prevented a severe reduction in glomerular filtration rate caused by ischemia/reperfusion or intra-arterial infusion of hydrogen peroxide [367]. This protective effect was ascribed, at least in part, to the induction of glomerular antioxidant enzymes, including Mn-SOD, catalase, and glutathione peroxidase.

As shown in Table 2, a number of defending molecules are inducible in response to proinflammatory or anti-inflammatory cytokines. Strictly controlled administration of cytokines might also achieve substantial enhancement of the glomerular self-defense.

### Pharmacological manipulation

Glucocorticoids are used for the first-line treatment of nephrotic syndrome, including minimal change disease in which oxidants are involved as pathogenic mediators. Ichikawa et al investigated the effect of glucocorticoid on the expression of antioxidant enzymes in the glomerulus. They found that, *in vitro*, glomerular endothelial and mesangial cells treated by methylprednisolone exhibited increased activity and expression of Mn-SOD and catalase [368]. They also showed that glomeruli isolated from rats treated with methylprednisolone had significantly higher activity of Mn-SOD, glutathione peroxidase, and catalase compared with control rats [393]. In PAN nephrosis, a model of minimal change disease, glucocorticoid-treated rats showed less impairment in renal function, lack of foot process fusion, and lower lipid peroxidation levels [393]. The therapeutic effect of glucocorticoids may be, in part, due to the enhancement of endogenous self-defense machinery that attenuates pathogenic action of ROI.

### Genetic manipulation

The progress in gene transfer technology has established the scientific basis for genetic engineering of somatic tissues. During the past years, several strategies have been developed to introduce exogenous genes into



the adult kidney [394, 395]. Using gene transfer technology, it should be feasible to reinforce self-defense of the glomerulus. Importantly, the gene transfer strategy is widely applicable; that is, a majority of defending molecules listed in Tables 1 and 2 are the potential targets for genetic manipulation.

*(1) Gene transfer approaches to the glomerulus.*

The introduction of foreign genes into the glomerulus is feasible via the renal circulation. Tomita et al reported an *in vivo* gene transfer approach using the hemagglutinating virus of Japan (HVJ)-liposome method [396]. In this approach, DNA, nuclear protein, and lipid are conjugated to form liposomes. Purified HVJ is inactivated by ultraviolet light and fused with the liposomes. By injection of the HVJ-liposome-coupled plasmid into the renal artery, expression of transgene is observed in glomerular cells, especially in mesangial cells [397]. Using this method, Badr et al showed that overexpression of 15-LO significantly improved glomerular filtration rate and proteinuria in anti-GBM nephritis in rats [abstract; Yura et al, *J Am Soc Nephrol* 6:890, 1995]. Another gene transfer approach was reported by Heikkila et al using a replication-deficient adenoviral vector [398]. In this method, pig kidneys are continuously perfused *in vivo* or *ex vivo* with an adenovirus that introduces a marker gene. Intense expression of the transgene product was detectable in glomerular cells, mainly in podocytes [399].

As an alternative approach, *ex vivo* gene transfer system has been developed by our group using the genetically engineered mesangial cell as a vector for gene delivery [155]. In this method, mesangial cells are cultured from isolated glomeruli, introduced with a foreign gene *in vitro*, and transferred back into the kidney via the renal circulation. Using  $\beta$ -galactosidase as a marker, we demonstrated efficient, site-specific gene delivery to the glomerulus and sustained transgene expression [155]. Using this approach, it is feasible to enhance glomerular self-defense. For example, the introduction of cDNA encoding IL-1ra conferred glomerular insensitiveness to IL-1 [400]. The transfer of a cDNA coding for active TGF- $\beta$ 1 reduced mitogenesis and cytokine responses of the glomerulus [154]. Fouqueray et al reported that intraglomerular transfer of a mesangial cell vector expressing human IL-10 resulted in a modest decrease in proteinuria in anti-GBM glomerulonephritis (abstract; Fouqueray et al, *J Am Soc Nephrol* 7:1698, 1996).

The approach to reinforcement of self-defense mechanisms has obvious requirements, the most important being that the defense systems should be bolstered only at a time point that is beneficial and that will not interfere with a protective inflammatory response. Toward achieving this goal, tight control over the transgene expression will be essential.

*(2) In vivo control of transgene activity.*

Many defending molecules are inducible in response

to a local inflammatory milieu. By combining gene transfer systems with inducible promoters, it is feasible to mimic the inducible property of endogenous defending molecules. Using the mesangial cell vector combined with the tetracycline-controlled transactivation system, we reported a reversible on-off system that achieves site-specific, *in vivo* control of transgene activity in the glomerulus [401]. By combining the mesangial cell vector with an "inflammation-responsive promoter," we further demonstrated automatic, *in vivo* induction of transgene expression in the glomerulus in response to local inflammatory stimuli [402]. These approaches would enable *in vivo* regulation of defender genes in which the transgene is required to be activated during inflammation and deactivated when the inflammation has subsided.

*(3) "Adoptive defense": Delivery of transgene products via the circulation.*


Increased defense potential of the glomerulus against complement-mediated injury may be achieved via adoptive transfer of GPI-anchored complement regulatory proteins. DAF and CD59 are GPI-anchored proteins that can be released from a cell membrane and reinserted into heterologous cell membranes without alteration of their biological function. Kooyman et al generated transgenic mice in which human DAF or CD59 was overexpressed on the surface of red blood cells. Using these mice, they demonstrated that functional DAF and CD59 are effectively transferred from red blood cells to the vascular endothelium [403]. Using this manipulation, McCurry et al showed that the intermembrane transfer of human DAF and CD59 effectively protected swine-to-primate cardiac xenografts from humoral injury [404]. This strategy may allow for the accumulation of complement regulatory proteins on the surface of glomerular endothelial cells without a direct gene transfer to the glomerulus.

Similar "adoptive defense" may be achieved in a different way. Recently, Zhu et al showed that a certain transgene product synthesized in the liver is effectively delivered to the glomerulus via the systemic circulation [405]. Isaka et al also reported that intramuscular transfer of a decorin cDNA led to accumulation of decorin protein in the glomerulus and attenuated matrix expansion in anti-Thy 1 glomerulonephritis.

## ACQUISITION BY THE GLOMERULUS OF "ANTI-INFLAMMATORY STATUS": A LESSON FROM ACUTE ANTI-THY 1 GLOMERULONEPHRITIS

As mentioned earlier, the concept of "glomerular self-defense" occurred to us via the finding that glomeruli in the regeneration phase of acute anti-Thy 1 glomerulonephritis (day 7) showed blunted responses to proinflammatory stimuli. As shown in Figure 2A, when normal





**Fig. 2. Blunted responses of nephritic glomeruli to stimuli.** (A) Blunted induction of metalloproteinase stromelysin in response to interleukin (IL)-1 $\beta$  in nephritic glomeruli. Glomeruli were isolated from normal rat kidneys (normal) and kidneys subjected to anti-Thy 1 glomerulonephritis at day 7 (anti-Thy 1 GN). The isolated glomeruli were stimulated by 10 ng/ml IL-1 $\beta$  for 24 hours in the presence of 1% fetal calf serum, and the induction of stromelysin was examined by Northern blot analysis. The expression of glyceraldehyde-3-phosphate dehydrogenase (GAPDH) is shown as a loading control. (B) Blunted responses of nephritic glomeruli to various stimuli. Isolated normal and nephritic glomeruli were stimulated by lipopolysaccharide (LPS; 1  $\mu$ g/ml), IL-1 $\beta$  (10 ng/ml), tumor necrosis factor- $\alpha$  (TNF- $\alpha$ ; 500 U/ml), platelet-derived growth factor (PDGF; 10 ng/ml), and 12-o-tetradecanoylphorbol-13-acetate (TPA; 50 ng/ml) for 24 hours. The expression of stromelysin, gelatinase B, and monocyte chemoattractant protein-1 (MCP-1) was examined by Northern blot analysis.


glomeruli were stimulated with the proinflammatory cytokine IL-1 $\beta$ , the expression of stromelysin was induced. Unexpectedly, however, the IL-1-triggered gene expression was blunted in the nephritic glomeruli. The blunted response was observed not only for stromelysin but also for other cytokine-inducible molecules, including gelatinase B and MCP-1. This insensitivity was consistently observed in the nephritic glomeruli triggered by other stimuli, including LPS and 12-o-tetradecanoylphorbol-13-acetate (TPA; Fig. 2B).

In the glomeruli seven days after the injection of the anti-Thy 1 antibody, mesangial cells are fully and preferentially activated, which suggests that defending molecules elaborated by mesangial cells may cause the blunted responses. At day 7 of anti-Thy 1 glomerulonephritis, macrophages are still present in the glomerulus [406]. Macrophage-derived factors might also participate in the reinforcement of glomerular self-defense. Based on these observations, several mechanisms can be postulated to explain the acquisition of the "anti-inflammatory status" by the nephritic glomeruli.

The first possibility is the elaboration of cytokine inhibitors such as IL-1ra. Although it has not been determined whether IL-1ra is expressed in anti-Thy 1 glomerulonephritis, monocytes and macrophages have the ability to produce IL-1ra abundantly. The insensitivity of the nephritic glomeruli to IL-1 may be ascribed to the elaborated IL-1ra.

The second possibility is that locally produced heparin-like molecules may inhibit activation of glomerular cells. Glomerular epithelial, endothelial, and mesangial cells have the ability to produce heparin-like substances [295, 296, 310, 311, 407]. In particular, mesangial cells produce HSPG in response to TGF- $\beta$  [312], which is abundantly elaborated in anti-Thy 1 glomerulonephritis [38]. *In vivo*, increased production of HSPG is observed in anti-Thy 1 glomerulonephritis [408]. HSPG inhibits mesangial expression of immediate early genes, including *c-fos* and *egr-1* [299, 300], which are involved in the induction of cytokine-inducible genes. The insensitivity of the nephritic glomeruli to stimuli might be due to local accumulation of heparin-like molecules.

The third possibility is that certain signaling pathways may be selectively suppressed via protein phosphatases. For example, a recent report showed that MKP-1 is up-regulated in the glomeruli of anti-Thy 1 glomerulonephritis [409]. MKP-1 suppresses activity of the MAP kinase family of molecules including ERK, JNK, and p38 MAP kinase [410, 411]. MAP kinases are generally involved in signaling pathways to transcriptional induction of various genes in response to inflammatory cytokines, growth factors, LPS, and pharmacological stimuli such as TPA [380]. The induction of protein phosphatases, including MKP-1, might be responsible for the attenuated responses of the nephritic glomeruli to external stimuli.



**Fig. 3. Inactivation of macrophages by bystander mesangial cells.** Normal rat macrophages (NR8383) were seeded on plastic or on confluent culture of rat mesangial cells (MC). After six hours, cells were stimulated by LPS (1  $\mu$ g/ml) for 14 hours. Then the macrophages were isolated from the cultures and subjected to Northern blot analyses of activation markers: TNF- $\alpha$ , inducible nitric oxide synthase (iNOS) and MCP-1.

An interesting finding is that the nephritic glomeruli containing many macrophages exhibit blunted responses to LPS and TPA, the strong activators for monocytes/macrophages. It indicates that, although macrophages are present, they become deactivated and thus insensitive to stimuli within the inflammatory milieu in which mesangial cells are activated. Cultured mesangial cells have the ability to deactivate macrophages [134, 135]. Figure 3 shows the LPS responses of rat macrophages cocultured with rat mesangial cells. The cocultured macrophages exhibit blunted expression of TNF- $\alpha$ , iNOS, and MCP-1 in response to LPS. Together with the fact that a number of defending molecules are produced by mesangial cells (Tables 1 and 2), this result raises the possibility that the mesangial cell is the pivotal defender of the glomerulus that functions against macrophage-mediated injury.

What kind of molecules do the nephritic glomeruli mobilize to inactivate macrophages? In this regard, macrophage-deactivating cytokines, including IL-4, IL-6, IL-10, IL-13 and TGF- $\beta$ , may play a role. Currently, information is limited regarding the expression of IL-4, IL-6, IL-10, and IL-13 in anti-Thy 1 nephritis, but it is well known that active TGF- $\beta$ 1 is abundantly produced in this nephritis model [38]. Several reports provided evidence that TGF- $\beta$ 1 inhibits production of IL-1 $\beta$ , TNF- $\alpha$ , and MCP-1 by activated macrophages [134, 144, 146]. TGF- $\beta$ 1 is a possible candidate that inactivates local macrophages.

Using a technique for *in vivo* macrophage transfer, it has been shown that glomerular cells in anti-Thy 1 glomerulonephritis are insensitive to the triggering action of activated macrophages (abstracts; Ogura et al, *ibid*; Sütö and Kitamura, *ibid*) [1, 2]. Cultured mesangial cells, glo-

merular endothelial cells, and isolated glomeruli treated with TGF- $\beta$ 1 exhibit depressed responses to IL-1 $\beta$  and TNF- $\alpha$  (abstract; Danoff and Jiang, *ibid*) [154, 159, 160]. TGF- $\beta$ 1-transfected mesangial cells show a blunted response to IL-1 $\beta$ , whereas mesangial cells expressing the mutant TGF- $\beta$  receptor show an enhanced response to IL-1, indicating that TGF- $\beta$ 1 functions as an autocrine inhibitor of the certain cytokine response in glomerular cells [1]. These data point to the potential of local TGF- $\beta$ 1 for suppression of acute inflammatory responses in the glomerulus via diminishing activity of both inflammatory macrophages and resident glomerular cells.

The anti-Thy 1 model (one shot model) mimics mesangial proliferative glomerulonephritis in humans, but the pathological changes are transient and reversible. After several weeks, the lesions spontaneously disappear without irreversible sclerosis or scarring. The anti-Thy 1 glomerulonephritis should therefore be considered as a model of glomerular repair, but not progressive glomerulosclerosis or scarring. During repair of the mesangiolytic glomerulonephritis, transient accumulation of ECM seems to be essential. This is because *in vivo* inactivation of TGF- $\beta$ 1 attenuates the accumulation of glomerular ECM, whereas it disturbs repair of mesangiolysis caused by anti-Thy 1 antibodies (Dr. Seiya Okuda, Kurume University School of Medicine, personal communication). It raises the possibility that TGF- $\beta$ 1 functions as an endogenous repair factor, rather than as a prosclerotic mediator, in this nephritis model. The "blackguard molecule" TGF- $\beta$ 1 should be rehabilitated under certain pathologic situations.

Currently, the contribution of NO and anti-inflammatory eicosanoids, the potential inactivators of leukocytes and glomerular cells, is undetermined. It is also unknown whether the insensitivity of nephritic glomeruli to stimuli is observed in other experimental glomerulonephritis. Extensive investigation is required to work out the molecular jigsaw of the glomerular self-defense system in individual human and experimental glomerular disease.

## PERSPECTIVE

The balance between external offending factors and internal defense machinery determines the fate of glomerular disease. A wide array of investigations have been pursued during the past years especially focusing on the offensive side of diseases. In contrast, only little attention has been paid to its defensive side. Information is still limited regarding what kind of endogenous molecules counteract complement, platelets, leukocytes and inflammatory mediators, and how their expression is controlled under pathophysiologic circumstances. Interestingly, some defending molecules are generated in response to their specific targets. For example, oxidants, IL-1, PDGF, TGF- $\beta$ , and complement trigger expression of their spe-

cific inhibitors—antioxidants, IL-1ra, SPARC, decorin, and complement regulatory proteins—in resident cells, respectively. Elucidation of this sophisticated, autoregulatory mechanism may be important for a better understanding of the glomerular defense system. One defending molecule often triggers the expression of other defenders. For example, TGF- $\beta$  induces IL-10, IL-1ra, SPARC, TIMP, HSPG, decorin, CD59, and CKIs (p15, p21, p27). IL-6 triggers the production of IL-1ra, sTNFR, PGE<sub>2</sub> and PGI<sub>2</sub>. IL-10 up-regulates IL-1ra and sTNFR. IL-4 and IL-13 stimulate IL-1ra, IL-1 decoy receptor and 15-LO expression. Heparin and HSPG induce endothelial NO, TFPI, HSPG, and potentiate function of C1 inhibitor, TGF- $\beta$ , and Mn-SOD. Intermolecular defense cascades and networks in the glomerulus are other important issues to be addressed in the future.

In addition to complement, platelets, leukocytes, and inflammatory mediators, vasoconstrictive substances are also considered pathogenic factors involved in glomerular injury [412]. Mechanical stress associated with local hypertension induces glomerular injury directly or indirectly [413]. In this regard, endogenous vasodepressors including NO and certain eicosanoids may form a line of self-defense [202, 314]. In this article, we have not dealt with hemodynamic-associated glomerular injury, but it should be another field in which self-defense is proposed to maintain glomerular integrity.

The establishment of effective therapeutic approaches to glomerular disease is still under way. Pharmacological inhibition of local pathogenic components is a possible approach. Alternatively, the reinforcement of endogenous defense mechanisms by pharmacologic and/or genetic manipulations may provide novel avenues to the treatment of glomerular disease. Efficient and realistic therapy would be feasible only if molecular events underlying glomerular diseases are fully elucidated from both the offensive and defensive sides.

## ACKNOWLEDGMENTS

This work was supported in part by grants from the Baxter Healthcare Corporation (Extramural Grant Program), the Wellcome Trust, and the National Kidney Research Fund to M. Kitamura.

Correspondence to Masanori Kitamura, M.D., Glomerular Bioengineering Unit, Department of Medicine, University College London Medical School, The Rayne Institute, 5 University Street, London WC1E 6JJ, United Kingdom.

E-mail: m.kitamura@medicine.ucl.ac.uk

## APPENDIX

Abbreviations used in this article are: AP-1, activator protein 1;  $\alpha$ -SMA,  $\alpha$ -smooth muscle actin; ATP/ADPase, adenosine triphosphatase and adenosine diphosphatase; AVP, arginine vasopressin; CDK, cyclin-dependent kinase; CKI, cyclin kinase inhibitor; CR1, complement receptor 1; Crry, complement receptor related gene y; DAF, decay accelerating factor; ECM, extracellular matrix; EC-SOD, extracellular superoxide dismutase; EGF, epidermal growth factor; ERK, extracellular

signal-regulated kinase; GBM, glomerular basement membrane; G-CSF, granulocyte colony stimulating factor; GLEPP1, glomerular epithelial protein 1; GM-CSF, granulocyte macrophage colony-stimulating factor; GPI, glycosyl-phosphatidylinositol; HSP, heat shock protein; HSPG, heparan sulfate proteoglycan; HVJ, hemagglutinating virus of Japan; ICAM-1, intercellular adhesion molecule 1; IL, interleukin; IL-1ra, IL-1 receptor antagonist; iNOS, inducible nitric oxide synthase; JNK, c-Jun N-terminal kinase; LIF, leukemia inhibitory factor; LO, lipoxygenase; LPS, lipopolysaccharide; LX, lipoxin; MAC, membrane attack complex; MAP kinase, mitogen-activated protein kinase; MCP, membrane cofactor protein; MCP-1, monocyte chemoattractant protein-1; MIF, macrophage migration inhibitory factor; MIP-1 $\alpha$ , macrophage inflammatory protein 1 $\alpha$ ; MIP-2, macrophage inflammatory protein-2; MKP-1, MAP kinase phosphatase 1; MMP, matrix metalloproteinase; NF- $\kappa$ B, nuclear factor- $\kappa$ B; NO, nitric oxide; p21, p21<sup>Cip1/Waf1/Sdi1/Cap20</sup>; p27, p27<sup>Kip1</sup>; PAN, puromycin aminonucleoside; PDGF, platelet-derived growth factor; PG, prostaglandin; PTP, protein-tyrosine phosphatase; PUMP, punctuated metalloproteinase; ROI, reactive oxygen intermediates; RPTP-BK, receptor-type PTP in the brain and kidney; SPARC, secreted protein acidic and rich in cysteine; sTNFR, soluble tumor necrosis factor receptor; TFPI, tissue factor pathway inhibitor; TGF- $\beta$ , transforming growth factor- $\beta$ ; Th2, T cell helper type 2; TIMP, tissue inhibitor of metalloproteinases; TNF- $\alpha$ , tumor necrosis factor- $\alpha$ ; t-PA, tissue-type plasminogen activator; TPA, 12-o-tetradecanoylphorbol-13-acetate; u-PA, urokinase-type plasminogen activator.

## REFERENCES

1. KITAMURA M: TGF- $\beta$  as an endogenous defender against macrophage-triggered stromelysin gene expression in the glomerulus. *J Immunol* 160:5163–5168, 1998
2. SUTÖ TS, FINE LG, SHIMIZU F, KITAMURA M: In vivo transfer of engineered macrophages into the glomerulus: Endogenous TGF- $\beta$ -mediated defence against macrophage-induced glomerular cell activation. *J Immunol* 159:2476–2483, 1997
3. SEDOR JR, NAKAZATO Y, KONIECZKOWSKI M: Interleukin-1 and the mesangial cell. *Kidney Int* 41:595–599, 1992
4. ATKINS RC: Interleukin-1 in crescentic glomerulonephritis. *Kidney Int* 48:576–586, 1995
5. AREND WP: Interleukin-1 receptor antagonist: A new member of the interleukin-1 family. *J Clin Invest* 88:1445–1451, 1991
6. DINARELLO CA, THOMPSON RC: Blocking IL-1: Interleukin 1 receptor antagonist in vivo and in vitro. *Immunol Today* 12:404–410, 1991
7. LAN HY, NIKOLIC-PATERSON DJ, ZARAMA M, VANNICE JL, ATKINS RC: Suppression of experimental crescentic glomerulonephritis by the interleukin-1 receptor antagonist. *Kidney Int* 43:479–485, 1993
8. TANG WW, FENG L, VANNICE JL, WILSON CB: Interleukin-1 receptor antagonist ameliorates experimental anti-glomerular basement membrane antibody-associated glomerulonephritis. *J Clin Invest* 93:273–279, 1994
9. NIKOLIC-PATERSON DJ, LAN HY, HILL PA, VANNICE JL, ATKINS RC: Suppression of experimental glomerulonephritis by the interleukin-1 receptor antagonist: Inhibition of intercellular adhesion molecule-1 expression. *J Am Soc Nephrol* 4:1695–1700, 1994
10. KARKAR AM, TAM FW, STEINKASSERER A, KURRLE R, LANGNER K, SCALLON BJ, MEAGER A, REES AJ: Modulation of antibody-mediated glomerular injury in vivo by IL-1ra, soluble IL-1 receptor, and soluble TNF receptor. *Kidney Int* 48:1738–1746, 1995
11. TAM FM, KARKAR AM, SMITH J, YOSHIMURA T, STEINKASSERER A, KURRLE R, LANGNER K, REES AJ: Differential expression of macrophage inflammatory protein-2 and monocyte chemoattractant protein-1 in experimental glomerulonephritis. *Kidney Int* 49:715–721, 1996
12. TESCH GH, LAN HY, ATKINS RC, NIKOLIC-PATERSON DJ: Role of interleukin-1 in mesangial cell proliferation and matrix deposition in experimental mesangioproliferative nephritis. *Am J Pathol* 151:141–150, 1997
13. LAN HY, NIKOLIC-PATERSON DJ, MU W, VANNICE JL, ATKINS RC: Interleukin-1 receptor antagonist halts the progression of established crescentic glomerulonephritis in the rat. *Kidney Int* 47:1303–1309, 1995



14. TAM FW, SMITH J, CASHMAN SJ, WANG Y, THOMPSON EM, REES AJ: Glomerular expression of interleukin-1 receptor antagonist and interleukin-1 $\beta$  genes in antibody-mediated glomerulonephritis. *Am J Pathol* 145:126–136, 1994
15. SMITH MF JR, EIDLEN D, AREND WP, GUTIERREZ-HARTMANN A: LPS-induced expression of the human IL-1 receptor antagonist gene is controlled by multiple interacting promoter elements. *J Immunol* 153:3584–3593, 1994
16. GABAY C, SMITH MF, EIDLEN D, AREND WP: Interleukin 1 receptor antagonist (IL-1ra) is an acute-phase protein. *J Clin Invest* 99:2930–2940, 1997
17. OHMORI Y, SMITH MF JR, HAMILTON TA: IL-4-induced expression of the IL-1 receptor antagonist gene is mediated by STAT6. *J Immunol* 157:2058–2065, 1996
18. WAHL SM, COSTA GL, CORCORAN M, WAHL LM, BERGER AE: Transforming growth factor- $\beta$  mediates IL-1-dependent induction of IL-1 receptor antagonist. *J Immunol* 150:3553–3560, 1993
19. BAUD L, ARDAILLOU R: Tumor necrosis factor- $\alpha$  in glomerular injury. *Kidney Int* 45(Suppl 45):S32–S36, 1994
20. MULLIGAN MS, JOHNSON KJ, TODD RF, ISSEKUTZ TB, MIYASAKA M, TAMATANI T, WAYNE-SMITH C, ANDERSON DC, WARD PA: Requirements for leukocyte adhesion molecules in nephrotoxic nephritis. *J Clin Invest* 91:577–587, 1993
21. HRUBY ZW, SHIROTA K, JOTHY S, LOWRY RP: Antiserum against tumor necrosis factor- $\alpha$  and a protease inhibitor reduce immune glomerular injury. *Kidney Int* 40:43–51, 1991
22. LAN HY, YANG N, METZ C, MU W, SONG Q, NIKOLIC-PATERSON DJ, BACHER M, BUCALA R, ATKINS RC: TNF- $\alpha$  up-regulates renal MIF expression in rat crescentic glomerulonephritis. *Mol Med* 3:136–144, 1997
23. SUZUKI J, TOMIZAWA S, ARAI H, SEKI Y, MARUYAMA K, KUROUME T: Purification of two types of TNF inhibitors in the urine of the patient with chronic glomerulonephritis. *Nephron* 66:386–390, 1994
24. LIEN E, LIABAKK NB, JOHNSEN AC, NONSTAD U, SUNDAN A, ESPEVIK T: Polymorphonuclear granulocytes enhance lipopolysaccharide-induced soluble p75 tumor necrosis factor receptor release from mononuclear cells. *Eur J Immunol* 25:2714–2717, 1995
25. JOHNSON RJ, FLOEGE J, COUSER WG, ALPERS CE: Role of platelet-derived growth factor in glomerular disease. *J Am Soc Nephrol* 4:119–128, 1993
26. JOHNSON RJ, RAINES EW, FLOEGE J, YOSHIMURA A, PRITZL P, ALPERS C, ROSS R: Inhibition of mesangial cell proliferation and matrix expansion in glomerulonephritis in the rat by antibody to platelet-derived growth factor. *J Exp Med* 175:1413–1416, 1992
27. RAINES EW, LANE TF, IRUELA-ARISPE ML, ROSS R, SAGE EH: The extracellular glycoprotein SPARC interacts with platelet-derived growth factor (PDGF)-AB and -BB and inhibits the binding of PDGF to its receptor. *Proc Natl Acad Sci USA* 89:1281–1285, 1992
28. CHANDRASEKHAR S, HARVEY AK, JOHNSON MG, BECKER GW: Osteonectin/SPARC is a product of articular chondrocytes/cartilage and is regulated by cytokines and growth factors. *Biochim Biophys Acta* 1221:7–14, 1994
29. FLOEGE J, JOHNSON RJ, ALPERS CE, FATEMI-NAINIE S, RICHARDSON CA, GORDON K, COUSER WG: Visceral glomerular epithelial cells can proliferate in vivo and synthesize platelet-derived growth factor B-chain. *Am J Pathol* 142:637–650, 1993
30. PICHLER RH, BASSUK JA, HUGO C, REED MJ, ENG E, GORDON KL, PIPPIN J, ALPERS CE, COUSER WG, SAGE EH, JOHNSON RJ: SPARC is expressed by mesangial cells in experimental mesangial proliferative nephritis and inhibits platelet-derived growth factor-mediated mesangial cell proliferation *in vitro*. *Am J Pathol* 148:1153–1167, 1996
31. BORDER WA, NOBLE NA: Transforming growth factor  $\beta$  in tissue fibrosis. *N Engl J Med* 331:1286–1292, 1994
32. ROBERTS AB, SPORN MB: The transforming growth factor- $\beta$ s, in *Peptide growth factors and their receptors*, edited by SPORN MB, ROBERTS AB, Berlin, Springer, 1990, pp 419–472
33. BORDER WA, OKUDA S, LANGUINO LR, SPORN MB, RUOSLAHTI E: Suppression of experimental glomerulonephritis by anti-serum against transforming growth factor  $\beta$ 1. *Nature* 346:371–374, 1990
34. AKAGI Y, ISAKA Y, ARAI M, KANEKO T, TAKENAKA M, MORIYAMA T, KANEDA Y, ANDO A, ORITA Y, KAMADA T, UEDA N, IMAI E: Inhibition of TGF- $\beta$ 1 expression by antisense oligonucleotides suppressed extracellular matrix accumulation in experimental glomerulonephritis. *Kidney Int* 50:148–155, 1996
35. BORDER WA, NOBLE NA, YAMAMOTO T, HARPER JR, YAMAGUCHI Y, PIERSCHBACHER MD, RUOSLAHTI E: Natural inhibitor of transforming growth factor- $\beta$  protects against scarring in experimental kidney disease. *Nature* 360:361–364, 1992
36. ISAKA Y, BRES DK, IKEGAYA K, KANEDA Y, IMAI E, NOBLE NA, BORDER WA: Gene therapy by skeletal muscle expression of decorin prevents fibrotic disease in rat kidney. *Nat Med* 2:418–423, 1996
37. BORDER WA, OKUDA S, LANGUINO LR, RUOSLAHTI E: Transforming growth factor- $\beta$  regulates production of proteoglycans by mesangial cells. *Kidney Int* 37:689–695, 1990
38. OKUDA S, LANGUINO LR, RUOSLAHTI E, BORDER WA: Elevated expression of transforming growth factor- $\beta$  and proteoglycan production in experimental glomerulonephritis: Possible role in expansion of the mesangial extracellular matrix. *J Clin Invest* 86:453–462, 1990
39. MATRISIAN LM: Metalloproteinases and their inhibitors in matrix remodeling. *Trends Genet* 6:121–125, 1990
40. NGUYEN HH, BARICOS WH, SHAH SV: Degradation of glomerular basement membrane by a neutral metalloproteinase(s) present in glomeruli isolated from normal rat kidney. *Biochem Biophys Res Commun* 141:898–903, 1986
41. LOVETT DH, JOHNSON RJ, MARTI H-P, MARTIN J, DAVIES M, COUSER WG: Structural characterization of the mesangial cell type IV collagenase and enhanced expression in a model of immune complex-mediated glomerulonephritis. *Am J Pathol* 141:85–98, 1992
42. EBIHARA I, NAKAMURA T, TOMINO Y, KOIDE H: Effect of a specific endothelin receptor A antagonist and an angiotensin-converting enzyme inhibitor on glomerular mRNA levels for extracellular matrix components, metalloproteinases (MMP) and a tissue inhibitor of MMP in aminonucleoside nephrosis. *Nephrol Dial Transplant* 12:1001–1006, 1997
43. DEMILLAN JJ, RIORDAN JW, COUSER WG, POLLOCK AS, LOVETT DH: Characterization of a glomerular epithelial cell metalloproteinase as matrix metalloproteinase-9 with enhanced expression in a model of membranous nephropathy. *J Clin Invest* 97:1094–1101, 1996
44. SUZUKI D, MIYAZAKI M, JINDE K, KOJI T, YAGAME M, ENDOH M, NOMOTO Y, SAKAI H: In situ hybridization studies of matrix metalloproteinase-3, tissue inhibitor of metalloproteinase-1 and type IV collagen in diabetic nephropathy. *Kidney Int* 52:111–119, 1997
45. MARTI HP, MCNEIL L, THOMAS G, DAVIES M, LOVETT DH: Molecular characterization of a low-molecular-mass matrix metalloproteinase secreted by glomerular mesangial cells as PUMP-1. *Biochem J* 285:899–905, 1992
46. BARICOS WH, SHAH SV: Proteolytic enzymes as mediators of glomerular injury. *Kidney Int* 40:161–173, 1991
47. SHAH SV, BARICOS WH, BASCI A: Degradation of human glomerular basement membrane by stimulated neutrophils: Activation of a metalloproteinase(s) by reactive oxygen metabolites. *J Clin Invest* 79:25–31, 1987
48. SHARMA R, SUZUKI K, NAGASE H, SAVIN VJ: Matrix metalloproteinase (stromelysin-1) increases the albumin permeability of isolated rat glomeruli. *J Lab Clin Med* 128:297–303, 1996
49. TURCK J, POLLOCK AS, LEE LK, MARTI HP, LOVETT DH: Matrix metalloproteinase 2 (gelatinase A) regulates glomerular mesangial cell proliferation and differentiation. *J Biol Chem* 271:15074–15083, 1996
50. STEINMANN-NIGGLI K, LUKES M, MARTI HP: Rat mesangial cells and matrix metalloproteinase inhibitor: Inhibition of 72-kD type IV collagenase (MMP-2) and of cell proliferation. *J Am Soc Nephrol* 8:395–405, 1997
51. KITAMURA M, SHIRASAWA T, MARUYAMA N: Gene transfer of metalloproteinase transin induces aberrant behavior of cultured mesangial cells. *Kidney Int* 45:1580–1586, 1994
52. WERB Z: ECM and cell surface proteolysis: Regulating cellular ecology. *Cell* 91:439–442, 1997



53. MARTIN J, KNOWLDEN J, DAVIES M, WILLIAMS JD: Identification and independent regulation of human mesangial cell metalloproteinases. *Kidney Int* 46:877-885, 1994
54. CAROME MA, STRIKER LJ, PETEN EP, MOORE J, YANG CW, STETLER-STEVENSON WG, STRIKER GE: Human glomeruli express TIMP-1 mRNA and TIMP-2 protein and mRNA. *Am J Physiol* 264:F923-F929, 1993
55. KITAMURA M, MARUYAMA N, MITARAI T, NAGASAWA R, YOKOO T, SAKAI O: Heparin selectively inhibits gene expression of matrix metalloproteinase transin in cultured mesangial cells. *Biochem Biophys Res Commun* 203:1333-1338, 1994
56. MARTI HP, LEE L, KASHGARIAN M, LOVETT DH: Transforming growth factor- $\beta$ 1 stimulates glomerular mesangial cell synthesis of the 72-kd type IV collagenase. *Am J Pathol* 144:82-94, 1994
57. KITAMURA M, KITAMURA A, MITARAI T, MARUYAMA N, NAGASAWA R, KAWAMURA T, YOSHIDA H, TAKAHASHI T, SAKAI O: Gene expression of metalloproteinase and its inhibitor in mesangial cells exposed to high glucose. *Biochem Biophys Res Commun* 185:1048-1054, 1992
58. SHANKLAND SJ, LY H, THAI K, SCHOLEY JW: Glomerular expression of tissue inhibitor of metalloproteinase (TIMP-1) in normal and diabetic rats. *J Am Soc Nephrol* 7:97-104, 1996
59. REDDY VY, ZHANG QY, WEISS SJ: Pericellular mobilization of the tissue-destructive cysteine proteinases, cathepsins B, L, and S, by human monocyte-derived macrophages. *Proc Natl Acad Sci USA* 92:3849-3853, 1995
60. KIMURA Y, YOKOI-HAYASHI K: Polymorphonuclear leukocyte lysosomal proteases, cathepsins B and D affect the fibrinolytic system in human umbilical vein endothelial cells. *Biochim Biophys Acta* 1310:1-4, 1997
61. BARICOS WH, CORTEZ SL, LE QC, ZHOU YW, DICARLO RM, O'CONNOR SE, SHAH SV: Glomerular basement membrane degradation by endogenous cysteine proteinases in isolated rat glomeruli. *Kidney Int* 38:395-401, 1990
62. BARICOS WH, O'CONNOR SE, CORTEZ SL, WU LT, SHAH SV: The cysteine proteinase inhibitor, E-64, reduces proteinuria in an experimental model of glomerulonephritis. *Biochem Biophys Res Commun* 155:1318-1323, 1988
63. BARICOS WH, CORTEZ SL, LE QC, WU LT, SHAW E, HANADA K, SHAH SV: Evidence suggesting a role for cathepsin L in an experimental model of glomerulonephritis. *Arch Biochem Biophys* 288:468-472, 1991
64. BRZIN J, POPOVIC T, TURK V, BORCHART U, MACHLEIDT W: Human cystatin, a new protein inhibitor of cysteine proteinases. *Biochem Biophys Res Commun* 118:103-109, 1984
65. ABRAHAMSON M, OLAFSSON I, PALSODTIR A, ULVSBACK M, LUNDWALL A, JENSSON O, GRUBB A: Structure and expression of the human cystatin C gene. *Biochem J* 268:287-294, 1990
66. TAVERA C, LEUNG-TACK J, PREVOT D, GENSAC MC, MARTINEZ J, FULCRAND P, COLLE A: Cystatin C secretion by rat mesangial cells: Autocrine loop for in vitro growth-promoting activity. *Biochem Biophys Res Commun* 182:1082-1088, 1992
67. QUIGG RJ: Glomerular injury induced by antibody and complement. *Semin Nephrol* 11:259-267, 1991
68. NANGAKU M, JOHNSON RJ, COUSER WG: Glomerulonephritis and complement regulatory proteins. *Exp Nephrol* 5:345-354, 1997
69. QUIGG RJ, NICHOLSON-WELLER A, CYBULSKY AV, BADALAMENTI J, SALANT DJ: Decay accelerating factor regulates complement activation on glomerular epithelial cells. *J Immunol* 142:877-882, 1989
70. ABE K, MIYAZAKI M, KOJI T, FURUSU A, OZONO Y, HARADA T, SAKAI H, NAKANE PK, KOHNO S: Expression of decay accelerating factor mRNA and complement C3 mRNA in human diseased kidney. *Kidney Int* 54:120-130, 1998
71. COSIO FG, SEDMAK DD, MAHAN JD, NAHMAN NS Jr: Localization of decay accelerating factor in normal and diseased kidneys. *Kidney Int* 36:100-107, 1989
72. SHIBATA T, COSIO FG, BIRMINGHAM DJ: Complement activation induces the expression of decay-accelerating factor on human mesangial cells. *J Immunol* 147:3901-3908, 1991
73. STORCK M, ABENDROTH D, PRESTEL R, PINO-CHAVEZ G, MULLER-HOKER J, WHITE DJ, HAMMER C: Morphology of hDAF (CD55) transgenic pig kidneys following ex-vivo hemoperfusion with human blood. *Transplantation* 63:304-310, 1997
74. QUIGG RJ, GALISHOFF ML, SNEED AE III, KIM D: Isolation and characterization of complement receptor type 1 from rat glomerular epithelial cells. *Kidney Int* 43:730-736, 1993
75. EMANCIPATOR SN, IIDA K, NUSSENZWEIG V, GALLO GR: Monoclonal antibodies to human complement receptor (CR1) detect defects in glomerular diseases. *Clin Immunol Immunopathol* 27:170-175, 1983
76. IIDA K, KOYAMA A, NAKAMURA H, INAGE H, NARITA M, TOJOYO S, KAMISATO J, FUJITA T, TAMURA N: Abnormal expression of complement receptor (CR1) in IgA nephritis: Increase in erythrocytes and loss on glomeruli in patients with impaired renal function. *Clin Immunol Immunopathol* 40:393-400, 1986
77. TEIXEIRA JE, COSTA RS, LACHMANN PJ, WURZNER R, BARBOSA JE: CR1 stump peptide and terminal complement complexes are found in the glomeruli of lupus nephritis patients. *Clin Exp Immunol* 105:497-503, 1996
78. COUSER WG, JOHNSON RJ, YOUNG BA, YEH CG, TOTH CA, RUDOLPH AR: The effects of soluble recombinant complement receptor 1 on complement-mediated experimental glomerulonephritis. *J Am Soc Nephrol* 5:1888-1894, 1995
79. NAKANISHI I, MOUTABARRIK A, HARA T, HATANAKA M, HAYASHI T, SYOUII T, OKADA N, KITAMURA E, TSUBAKIHARA Y, MATSUMOTO M, SEYA T: Identification and characterization of membrane cofactor protein (CD46) in the human kidneys. *Eur J Immunol* 24:1529-1535, 1994
80. COSIO FG, SHIBATA T, ROVIN BH, BIRMINGHAM DJ: Effects of complement activation products on the synthesis of decay accelerating factor and membrane cofactor protein by human mesangial cells. *Kidney Int* 46:986-989, 1994
81. ENDOH M, YAMASHINA M, OHI H, FUNAHASHI K, IKUNO T, YASUGI T, ATKINSON JP, OKADA H: Immunohistochemical demonstration of membrane cofactor protein (MCP) of complement in normal and diseased kidney tissues. *Clin Exp Immunol* 94:182-188, 1993
82. QUIGG RJ, HOLERS VM, MORGAN BP, SNEED AE III: Crry and CD59 regulate complement in rat glomerular epithelial cells and are inhibited by the nephritogenic antibody of passive Heymann nephritis. *J Immunol* 154:3437-3443, 1995
83. QUIGG RJ, MORGAN BP, HOLERS VM, ADLER S, SNEED AE III, LO CF: Complement regulation in the rat glomerulus: Crry and CD59 regulate complement in glomerular mesangial and endothelial cells. *Kidney Int* 48:412-421, 1995
84. MOUTABARRIK A, NAKANISHI I, MATSUMOTO M, ZAID D, SEYA T: Human glomerular epithelial cells synthesize and secrete the third component of complement. *Nephron* 70:55-61, 1995
85. SHIBATA T, KOHSAKA T: Effects of complement activation on the expression of CD59 by human mesangial cells. *J Immunol* 155:403-409, 1995
86. NANGAKU M, MEEK RL, PIPPIN J, GORDON KL, MORGAN BP, JOHNSON RJ, COUSER WG: Transfected CD59 protects mesangial cells from injury induced by antibody and complement. *Kidney Int* 50:257-266, 1996
87. TAMAI H, MATSUO S, FUKATSU A, NISHIKAWA K, SAKAMOTO N, YOSHIOKA K, OKADA N, OKADA H: Localization of 20-kD homologous restriction factor (HRF20) in diseased human glomeruli: An immunofluorescence study. *Clin Exp Immunol* 84:256-262, 1991
88. LEHTO T, HONKANEN E, TEPPA AM, MERI S: Urinary excretion of protectin (CD59), complement SC5b-9 and cytokines in membranous glomerulonephritis. *Kidney Int* 47:1403-1411, 1995
89. MATSUO S, NISHIKAGE H, YOSHIDA F, NOMURA A, PIDDLESSEN SJ, MORGAN BP: Role of CD59 in experimental glomerulonephritis in rats. *Kidney Int* 46:191-200, 1994
90. NANGAKU M, ALPERS CE, PIPPIN J, SHANKLAND SJ, KUROKAWA K, ADLER S, MORGAN BP, JOHNSON RJ, COUSER WG: CD59 protects glomerular endothelial cells from immune-mediated thrombotic microangiopathy in rats. *J Am Soc Nephrol* 9:590-597, 1998
91. NANGAKU M, QUIGG RJ, SHANKLAND SJ, OKADA N, JOHNSON RJ, COUSER WG: Overexpression of Crry protects mesangial cells from complement-mediated injury. *J Am Soc Nephrol* 8:223-233, 1997
92. MATSUO S, ICHIDA S, TAKIZAWA H, OKADA N, BARANYI L, IGUCHI A, MORGAN BP, OKADA H: In vivo effects of monoclonal antibody

- ies that functionally inhibit complement regulatory proteins in rats. *J Exp Med* 180:1619–1627, 1994
93. NISHIKAGE H, BARANYI L, OKADA H, OKADA N, ISOBE K, NOMURA A, YOSHIDA F, MATSUO S: The role of a complement regulatory protein in rat mesangial glomerulonephritis. *J Am Soc Nephrol* 6:234–241, 1995
  94. LAUFER J, OREN R, FARZAM N, GOLDBERG I, PASSWELL J: Differential cytokine regulation of complement proteins in human glomerular epithelial cells. *Nephron* 76:276–283, 1997
  95. HOGASEN K, JANSEN JH, MOLLNES TE, HOVDENES J, HARBOE M: Hereditary porcine membranoproliferative glomerulonephritis type II is caused by factor H deficiency. *J Clin Invest* 95:1054–1061, 1995
  96. FIEN CA, KUIJPER EJ, TE BULTE M, VAN DE HEUVEL MM, HOLDRIET AC, SIM RB, DAHA MR, DANKERT J: Heterozygous and homozygous factor H deficiency states in a Dutch family. *Clin Exp Immunol* 105:511–516, 1996
  97. TIMMERMAN JJ, VAN DER WOUDE FJ, VAN GILSWIJK-JANSSEN DJ, VERWEIJ CL, VAN ES LA, DAHA MR: Differential expression of complement components in human fetal and adult kidneys. *Kidney Int* 49:730–740, 1996
  98. AN DEL DOBBELSTEEN ME, VERHASSELT V, KAASHOEK JG, TIMMERMAN JJ, SCHROEIJERS WE, VERWEIJ-CL, VAN DER WOUDE FJ, VAN ES LA, DAHA MR: Regulation of C3 and factor H synthesis of human glomerular mesangial cells by IL-1 and interferon- $\gamma$ . *Clin Exp Immunol* 95:173–180, 1994
  99. BARIETY J, HINGLAIS N, BHAKDI S, MANDET C, ROUCHON M, KAZATCHKINE MD: Immunohistochemical study of complement S protein (Vitronectin) in normal and diseased human kidneys: Relationship to neoantigens of the C5b-9 terminal complex. *Clin Exp Immunol* 75:76–81, 1989
  100. OKADA M, YOSHIOKA K, TAKEMURA T, AKANO N, AYA N, MURAKAMI K, MAKI S: Immunohistochemical localization of C3d fragment of complement and S-protein (vitronectin) in normal and diseased human kidneys: Association with the C5b-9 complex and vitronectin receptor. *Virchows Arch A Pathol Anat Histopathol* 422:367–373, 1993
  101. MURPHY BF, KIRSZBAUM L, WALKER ID, D'APICE AJF: SP-40,40 a newly identified normal human serum protein found in the SC5b-9 complex of complement and in the immune deposits of glomerulonephritis. *J Clin Invest* 81:1858–1864, 1988
  102. ROSENBERG ME, SILKENSEN J: Clusterin: Physiologic and pathophysiologic considerations. *Int J Biochem Cell Biol* 27:633–645, 1995
  103. FRENCH LE, TSCHOPP J, SCHIFFERLI JA: Clusterin in renal tissue: Preferential localization with the terminal complement complex and immunoglobulin deposits in glomeruli. *Clin Exp Immunol* 88:389–393, 1992
  104. MURPHY BF, DAVIES DJ, MORROW W, D'APICE AJF: Localization of terminal complement components, S-protein and SP-40,40 in renal biopsies. *Pathology* 21:275–278, 1989
  105. EDDY AA, FRITZ IB: Localization of clusterin in the epimembranous deposits of passive Heymann nephritis. *Kidney Int* 39:247–252, 1991
  106. LAPING NJ, OLSON BA, SHORT B, ALBRIGHTSON CR: Thrombin increases clusterin mRNA in glomerular epithelial and mesangial cells. *J Am Soc Nephrol* 8:906–914, 1997
  107. SAUNDERS JR, AMINIAN A, McRAE JL, O'FARRELL KA, ADAM WR, MURPHY BF: Clusterin depletion enhances immune glomerular injury in the isolated perfused kidney. *Kidney Int* 45:817–827, 1994
  108. HALLIWELL B, GUTTERIDGE JMC: The antioxidants of human extracellular fluids. *Arch Biochem Biophys* 280:1–8, 1990
  109. MARKLUND SL: Extracellular superoxide dismutase in human tissues and human cell lines. *J Clin Invest* 74:1398–1403, 1984
  110. BAKKER WW, POELSTRA K, HARDONK MJ: Relevance of adenosine nucleotidases in glomerular filtration barrier. *Nephron* 64:338–342, 1993
  111. POELSTRA K, HEYENEN ER, BALLER JF, HARDONK MJ, BAKKER WW: Modulation of anti-Thy1 nephritis in the rat by adenosine nucleotides: Evidence for an anti-inflammatory role for nucleotidases. *Lab Invest* 66:555–563, 1992
  112. BAKKER WW, WILLINK EJ, DONGA J, HULSTAERT CE, HARDONK MJ: Antithrombotic activity of glomerular adenosine diphosphatase in the glomerular basement membrane of the rat kidney. *J Lab Clin Med* 109:171–177, 1987
  113. BAKKER WW, POELSTRA K, TIMMERMAN W, HARDONK MJ, KOITER TR, SCHULING GA: Experimental endotoxemia in pregnancy: In situ glomerular microthrombus formation associated with impaired glomerular adenosine diphosphatase activity. *J Lab Clin Med* 114:531–537, 1989
  114. POELSTRA K, HARDONK MJ, KOUDESTAAL J, BAKKER WW: Intraglomerular platelet aggregation and experimental glomerulonephritis. *Kidney Int* 37:1500–1508, 1990
  115. POELSTRA K, BALLER JF, HARDONK MJ, BAKKER WW: Intraglomerular thrombotic tendency and glomerular ADPase: Unilateral impairment of ADPase elicits a proaggregatory microenvironment in experimental glomerulonephritis. *Lab Invest* 64:520–526, 1991
  116. POELSTRA K, BALLER JF, HARDONK MJ, BAKKER WW: Demonstration of antithrombotic activity of glomerular adenosine diphosphatase. *Blood* 78:141–148, 1991
  117. HOOKE DH, GEE DC, ATKINS RC: Leukocytes analysis using monoclonal antibodies in human glomerulonephritis. *Kidney Int* 31:964–972, 1987
  118. MAIN IW, NIKOLIC-PATERSON DJ, ATKINS RC: T cells and macrophages and their role in renal injury. *Semin Nephrol* 12:395–407, 1992
  119. KITAMURA M, SÜTÖ TS: TGF- $\beta$  and glomerulonephritis: Anti-inflammatory versus prosclerotic actions. *Nephrol Dial Transplant* 12:669–679, 1997
  120. WAHL SM, MCCARTNEY-FRANCIS N, MERGENHAGEN SE: Inflammatory and immunomodulatory roles of TGF- $\beta$ . *Immunol Today* 10:258–261, 1989
  121. BRANDES ME, ALLEN JB, OGAWA Y, WAHL SM: Transforming growth factor- $\beta$ 1 suppresses acute and chronic arthritis in experimental animals. *J Clin Invest* 87:1108–1113, 1991
  122. KURUVILLA AP, SHAH R, HOCHWALD GM, LIGGITT HD, PALLADINO MA, THORBECKE GJ: Protective effect of transforming growth factor- $\beta$ 1 on experimental autoimmune diseases in mice. *Proc Natl Acad Sci USA* 88:2918–2921, 1991
  123. DANG H, GEISER AG, LETTERIO JJ, NAKABAYASHI T, KONG L, FERNANDES G, TALAL N: SLE-like autoantibodies and Sjögren syndrome-like lymphoproliferation in TGF- $\beta$  knockout mice. *J Immunol* 155:3205–3212, 1995
  124. RAZ E, WATANABE A, BAIRD SM, EISENBERG RA, PARR TB, LOTZ M, KIPPS T, CARSON DA: Systemic immunological effects of cytokine genes injected into skeletal muscle. *Proc Natl Acad Sci USA* 90:4523–4527, 1993
  125. SHULL MM, ORMSBY I, KIER AB, PAWLOWSKI S, DIEBOLD RJ, YIN M, ALLEN R, SIDMAN C, PROETZEL G, CALVIN D, ANNUZIATA N, DOETSCHMAN T: Targeted disruption of the mouse transforming growth factor- $\beta$ 1 null mutation in mice causes excessive inflammatory response and early death. *Proc Natl Acad Sci USA* 90:770–774, 1993
  126. KULKARNI AB, HUH CG, BECKER D, GEISER A, LYGT M, FLANDERS KC, ROBERTS AB, SPORN MB, WARD JM, KARLSSON S: Transforming growth factor- $\beta$ 1 null mutation in mice causes excessive inflammatory response and early death. *Proc Natl Acad Sci USA* 90:770–774, 1993
  127. CHRIST M, MCCARTNEY-FRANCIS NL, KULKARNI AB, WARD JM, MIZEL DE, MACKALL CL, GRESS RE, HINES KL, TIAN H, KARLSSON S, WAHL SM: Immune dysregulation in TGF- $\beta$ 1-deficient mice. *J Immunol* 153:1936–1946, 1994
  128. MACKAY K, KONDAIAH P, DANIELPOUR D, AUSTIN HA III, BROWN PD: Expression of transforming growth factor- $\beta$ 1 and  $\beta$ 2 in rat glomeruli. *Kidney Int* 38:1095–1100, 1990
  129. ANDO T, OKUDA S, TAMAKI K, YOSHITOMI K, FUJISHIMA M: Localization of transforming growth factor- $\beta$  and latent transforming growth factor- $\beta$  binding protein in rat kidney. *Kidney Int* 47:733–739, 1995
  130. YAMAMOTO T, NOBLE NA, COHEN AH, NAST CC, HISHIDA A, GOLD LI, BORDER WA: Expression of transforming growth factor- $\beta$  isoforms in human glomerular diseases. *Kidney Int* 49:461–469, 1996
  131. YAMAMOTO T, NOBLE NA, MILLER DE, BORDER WA: Sustained

- expression of TGF- $\beta$ 1 underlies development of progressive kidney fibrosis. *Kidney Int* 45:916–927, 1994
132. VAN DET NF, VERHAGEN NA, TAMSMA JT, BERDEN JH, BRUIJN JA, DAHA MR, VAN DER WOUDE FJ: Regulation of glomerular epithelial cell production of fibronectin and transforming growth factor- $\beta$  by high glucose, not by angiotensin II. *Diabetes* 46:834–840, 1997
  133. KANAME S, UCHIDA S, OGATA E, KUROKAWA K: Autocrine secretion of transforming growth factor- $\beta$  in cultured rat mesangial cells. *Kidney Int* 42:1319–1327, 1992
  134. KITAMURA M, SÜTÖ TS, YOKOO T, SHIMIZU F, FINE LG: Transforming growth factor- $\beta$ 1 is the predominant paracrine inhibitor of macrophage cytokine synthesis produced by glomerular mesangial cells. *J Immunol* 156:2964–2971, 1996
  135. SÜTÖ TS, FINE LG, KITAMURA M: Mesangial cell-derived transforming growth factor- $\beta$ 1 reduces macrophage adhesiveness with consequent deactivation. *Kidney Int* 50:445–452, 1996
  136. KAGAMI S, BORDER WA, MILLER DE, NOBLE NA: Angiotensin II stimulates extracellular matrix protein synthesis through induction of transforming growth factor- $\beta$  expression in rat glomerular mesangial cells. *J Clin Invest* 93:2431–2437, 1994
  137. WOLF G, ZIYADEH FN, STAHL RA: Atrial natriuretic peptide stimulates the expression of transforming growth factor- $\beta$  in cultured murine mesangial cells: Relationship to suppression of proliferation. *J Am Soc Nephrol* 6:224–233, 1995
  138. YAMABE H, OSAWA H, INUMA H, KAIZUKA M, TAMURA N, TSUNODA S, BABA Y, SHIRATO K, ONODERA K: Thrombin stimulates production of transforming growth factor- $\beta$  by cultured human mesangial cells. *Nephrol Dial Transplant* 12:438–442, 1997
  139. STUDER RK, DERUBERTIS FR, CRAVEN PA: Nitric oxide suppresses increases in mesangial cell protein kinase C, transforming growth factor  $\beta$ , and fibronectin synthesis induced by thromboxane. *J Am Soc Nephrol* 7:999–1005, 1996
  140. DI PAOLO S, GESUALDO L, RANIERI E, GRANDALIANO G, SCHENA FP: High glucose concentration induces the overexpression of transforming growth factor- $\beta$  through the activation of a platelet-derived growth factor loop in human mesangial cells. *Am J Pathol* 149:2095–2106, 1996
  141. DING G, VAN GOOR H, RICARDO SD, ORLOWSKI JM, DIAMOND JR: Oxidized LDL stimulates the expression of TGF- $\beta$  and fibronectin in human glomerular epithelial cells. *Kidney Int* 51:147–154, 1997
  142. LOPEZ-ARMADA MJ, GOMEZ-GUERRERO C, EGIDO J: Receptors for immune complexes activate gene expression and synthesis of matrix proteins in cultured rat and human mesangial cells: Role of TGF- $\beta$ . *J Immunol* 157:2136–2142, 1996
  143. RISER BL, CORTES P, HEILIG C, GRONDIN J, LADSON-WOFFORD S, PATTERSON D, NARINS RG: Cyclic stretching force selectively up-regulates transforming growth factor- $\beta$  isoforms in cultured rat mesangial cells. *Am J Pathol* 148:1915–1923, 1996
  144. ESPEVIK T, FIGARI IS, SHALABY MR, LACKIDES GA, LEWIS GD, SHEPARD HM, PALLADINO MA: Inhibition of cytokine production by cyclosporin A and transforming growth factor  $\beta$ . *J Exp Med* 166:571–576, 1987
  145. BOGDAN C, PAIK J, VODOCOTZ Y, NATHAN C: Contrasting mechanisms for suppression of macrophage cytokine release by transforming growth factor- $\beta$  and interleukin-10. *J Biol Chem* 267:23301–23308, 1992
  146. KITAMURA M: Identification of an inhibitor targeting macrophage production of monocyte chemoattractant protein-1 as transforming growth factor- $\beta$ 1. *J Immunol* 159:1404–1411, 1997
  147. SHAH SV: The role of reactive oxygen metabolites in glomerular disease. *Ann Rev Physiol* 57:245–262, 1995
  148. CATTELL V, COOK HT: Nitric oxide: Role in the physiology and pathology of the glomerulus. *Exp Nephrol* 1:265–280, 1993
  149. TSUNAWAKI S, SPORN MB, DING A, NATHAN C: Deactivation of macrophages by transforming growth factor- $\beta$ . *Nature* 334:260–262, 1988
  150. DING A, NATHAN CF, GRAYCAR J, DERYNCK R, STUEHR DJ, SRIMAL S: Macrophage deactivating factor and transforming growth factor- $\beta$ 1, - $\beta$ 2, - $\beta$ 3 inhibit induction of macrophage nitrogen oxide synthesis by IFN- $\gamma$ . *J Immunol* 145:940–944, 1990
  151. JAFFER F, SAUNDERS C, SCHULTZ P, THROCKMORTON D, WEINSHILL E, ABBODD HE: Regulation of mesangial cell growth by polypeptide mitogens: Inhibitory role of transforming growth factor- $\beta$ . *Am J Pathol* 135:261–269, 1989
  152. ADLER S, CHEN X, ENG B: Control of rat glomerular epithelial cell growth in vitro. *Kidney Int* 37:1048–1054, 1990
  153. CHOI ME, BALLERMANN BJ: Inhibition of capillary morphogenesis and associated apoptosis by dominant negative mutant transforming growth factor- $\beta$  receptors. *J Biol Chem* 270:21144–21150, 1995
  154. KITAMURA M, BURTON S, ENGLISH J, KAWACHI H, FINE LG: Transfer of a mutated gene encoding active transforming growth factor- $\beta$ 1 suppresses mitogenesis and IL-1 response in the glomerulus. *Kidney Int* 48:1747–1757, 1995
  155. KITAMURA M, TAYLOR S, UNWIN R, BURTON S, SHIMIZU F, FINE LG: Gene transfer into the rat renal glomerulus via a mesangial cell vector: Site-specific delivery, in situ amplification, and sustained expression of an exogenous gene in vivo. *J Clin Invest* 94:497–505, 1994
  156. REDINI F, MAUVIEL A, PRONOST S, LOYAU G, PUJOL J-P: Transforming growth factor- $\beta$  exerts opposite effects from interleukin-1 $\beta$  on cultured rabbit articular chondrocytes through reduction of interleukin-1 receptor expression. *Arthritis Rheum* 36:44–50, 1993
  157. VAN BEUNINGEN HM, VAN DER KRAAN PM, ARNTZ OJ, VAN DER BERG WB: Protection from interleukin 1 induced destruction of articular cartilage by transforming growth factor- $\beta$ : Studies in anatomically intact cartilage in vitro and in vivo. *Ann Rheum Dis* 52:185–191, 1993
  158. DUBOIS CM, RUSCETTI FW, PALASZYNSKI EW, FALK LA, OPPENHEIM JJ, KELLER JR: Transforming growth factor  $\beta$  is a potent inhibitor of IL-1 receptor expression: Proposed mechanism of inhibition on IL-1 action. *J Exp Med* 172:737–744, 1990
  159. PFEILSCHIFTER J, VOSBECK K: Transforming growth factor  $\beta$ 2 inhibits interleukin 1 $\beta$ - and tumour necrosis factor  $\alpha$ -induction of nitric oxide synthase in rat renal mesangial cells. *Biochem Biophys Res Commun* 175:372–379, 1991
  160. VERVOORDELDONK MJ, SCHALKWIJK CG, PFEILSCHIFTER J, VAN DEN BOSCH H: Effects of dexamethasone and transforming growth factor- $\beta$ 2 on group II phospholipase A<sub>2</sub> mRNA and activity levels in interleukin 1 $\beta$  and forskolin-stimulated mesangial cells. *Biochem J* 315:435–441, 1996
  161. KITAMURA M, SÜTÖ TS: Transfer of genetically engineered macrophages into the glomerulus. *Kidney Int* 51:1274–1279, 1997
  162. LAKKIS FG, BADDOURA FK, CRUET EN, PAREKH KR, FUKUNAGA M, MUNGER KA: Anti-inflammatory lymphokine mRNA expression in antibody-induced glomerulonephritis. *Kidney Int* 49:117–126, 1996
  163. OKADA H, KONISHI K, NAKAZATO Y, KANNO Y, SUZUKI H, SAKAGUCHI H, SARUTA T: Interleukin-4 expression in mesangial proliferative glomerulonephritis. *Am J Kidney Dis* 23:242–246, 1994
  164. FURUSU A, MIYAZAKI M, KOJI T, ABE K, OZONO Y, HARADA T, NAKANE PK, HARA K, KOHNO S: Involvement of IL-4 in human glomerulonephritis: An in situ hybridization study of IL-4 mRNA and IL-4 receptor mRNA. *J Am Soc Nephrol* 8:730–741, 1997
  165. KLUTH DC, REES AJ: Inhibiting inflammatory cytokines. *Semin Nephrol* 16:576–582, 1996
  166. TIPPING PG, KITCHING AR, HUANG XR, MUTCH DA, HOLDSWORTH SR: Immune modulation with interleukin-4 and interleukin-10 prevents crescent formation and glomerular injury in experimental glomerulonephritis. *Eur J Immunol* 27:530–537, 1997
  167. KITCHING AR, TIPPING PG, MUTCH DA, HUANG XR, HOLDSWORTH SR: Interleukin-4 deficiency enhances Th1 responses and crescentic glomerulonephritis in mice. *Kidney Int* 53:112–118, 1998
  168. ERB KJ, RUGER B, VON BREVERN M, RYFFEL B, SCHIMPL A, RIVETT K: Constitutive expression of interleukin-4 in vivo causes autoimmune-type disorders in mice. *J Exp Med* 185:329–339, 1997
  169. SCHORLEMMER HU, DICKNEITE G, KANZY EJ, ENSLE KH: Modulation of the immunoglobulin dysregulation in GvH- and SLE-like diseases by the murine IL-4 receptor (IL-4-R). *Inflamm Res* 44(Suppl 2):S194–S196, 1995
  170. NAKAZATO Y, OKADA H, SATO A, IWAITA Y, HAYASHIDA T, HAYASHI M, SUZUKI H, SARUTA T: Interleukin 4 downregulates cell growth and prostaglandin release of human mesangial cells. *Biochem Biophys Res Commun* 197:486–493, 1993



171. SAVAGE CO, BROOKS CJ, ADU D, RICHARDS G, HOWIE AJ: Cell adhesion molecule expression within human glomerular and kidney organ culture. *J Pathol* 181:111–115, 1997
172. MOUTABARRIK A, NAKANISHI I, ISHIBASHI M: Interleukin-6 and interleukin-6 receptor are expressed by cultured glomerular epithelial cells. *Scand J Immunol* 40:181–186, 1994
173. HORII Y, MURAGUCHI A, IWANO M, MATSUDA T, HIRAYAMA T, YAMADA H, FUJII Y, DOHI K, ISHIKAWA H, OHMOTO Y, YOSHIZAKI K, HIRANO T, KISHIMOTO T: Involvement of IL-6 in mesangial proliferative glomerulonephritis. *J Immunol* 143:3949–3955, 1989
174. ZOJA C, WANG JM, BETTONI S, SIRONI M, RENZI D, CHIAFFARINO F, ABBOUD HE, VAN-DAMME J, MANTOVANI A, REMUZZI G: Interleukin-1 $\beta$  and tumor necrosis factor- $\alpha$  induce gene expression and production of leukocyte chemotactic factors, colony-stimulating factors, and interleukin-6 in human mesangial cells. *Am J Pathol* 138:991–1003, 1991
175. ZIESCHE R, ROTH M, PAPAKONSTANTINOU E, NAUCK M, HÖRL WH, KASHGARIAN M, BLOCK LH: A granulocyte inhibitory protein overexpressed in chronic renal disease regulates expression of interleukin 6 and interleukin 8. *Proc Natl Acad Sci USA* 91:301–305, 1994
176. MORIYAMA T, FUJIBAYASHI M, FUJIWARA Y, KANEKO T, XIA C, IMAI E, KAMADA T, ANDO A, UEDA N: Angiotensin II stimulates interleukin-6 release from cultured mouse mesangial cells. *J Am Soc Nephrol* 6:95–101, 1995
177. VAN DEN DOBBELSTEEN ME, VAN DER WOUDE FJ, SCHROEIJERS WE, VAN ES LA, DAHA MR: Soluble aggregates of IgG and immune complexes enhance IL-6 production by renal mesangial cells. *Kidney Int* 43:544–553, 1993
178. FUKATSU A, MATSUO S, TAMAI H, SAKAMOTO N, MATSUDA T, HIRANO T: Distribution of interleukin-6 in normal and diseased human kidney. *Lab Invest* 65:61–66, 1991
179. CHEN WP, LIN CY: Augmented expression of interleukin-6 and interleukin-1 genes in the mesangium of IgM mesangial nephropathy. *Nephron* 68:10–19, 1994
180. SUEMATSU S, MATSUDA T, AOZASA K, AKIRA S, NAKANO N, OHNO S, MIYAZAKI J, YAMAMURA K, HIRANO T, KISHIMOTO T: IgG1 plasmacytosis in interleukin 6 transgenic mice. *Proc Natl Acad Sci USA* 86:7547–7551, 1989
181. HARTNER A, GOPPELT-STRUEBE M, HOCKE GM, STERZEL RB: Differential regulation of chemokines by leukemia inhibitory factor, interleukin-6 and oncostatin M. *Kidney Int* 51:1754–1760, 1997
182. RYFFEL B, CAR BD, GUNN H, ROMAN D, HIESTAND P, MIHATSCH MJ: Interleukin-6 exacerbates glomerulonephritis in (NZB x NZW) F1 mice. *Am J Pathol* 144:927–937, 1994
183. KIBERD BA: Interleukin-6 receptor blockage ameliorates murine lupus nephritis. *J Am Soc Nephrol* 4:58–61, 1993
184. FLOEGE J, ENG E, YOUNG BA, ALPERS CE, BARRETT TB, BOWEN-POPE DF, JOHNSON RJ: Infusion of platelet-derived growth factor or basic fibroblast growth factor induces selective glomerular mesangial cell proliferation and matrix accumulation in rats. *J Clin Invest* 92:2952–2962, 1993
185. EITNER F, WESTERHUIS R, BURG M, WEINHOLD B, GRONE HJ, OSTENDORF T, RUTHER U, KOCH KM, REES AJ, FLOEGE J: Role of interleukin-6 in mediating mesangial cell proliferation and matrix production in vivo. *Kidney Int* 51:69–78, 1997
186. IKEDA M, IKEDA U, OHARA T, KUSANO E, KANO S: Recombinant interleukin-6 inhibits the growth of rat mesangial cells in culture. *Am J Pathol* 141:327–334, 1992
187. MATSELL DG, GABER LW, MALIK KU: Cytokine stimulation of prostaglandin production inhibits the proliferation of serum-stimulated mesangial cells. *Kidney Int* 45:159–165, 1994
188. KARKAR AM, TAM FW, PROUDFOOT AE, MEAGER A, REES AJ: Modulation of antibody-mediated glomerular injury in vivo by interleukin-6. *Kidney Int* 44:967–973, 1993
189. XING Z, GAULDIE J, COX G, BAUMANN H, JORDANA M, LEI X-F, ACHONG MK: IL-6 is an anti-inflammatory cytokine required for controlling local and systemic acute inflammatory responses. *J Clin Invest* 101:311–320, 1998
190. FIORENTINO DF, ZLOTNIK A, MOSMANN TR, HOWARD MH, O'GARRA A: IL-10 inhibits cytokine production by activated macrophages. *J Immunol* 147:3815–3822, 1991
191. DE WAAL-MALEFYT R, ABRAMS J, BENNETT B, FIGDOR C, DE VRIES J: IL-10 inhibits cytokine synthesis by human monocytes: An auto-regulatory role of IL-10 produced by monocytes. *J Exp Med* 174:1209–1220, 1991
192. BOGDAN C, VODOVOTZ Y, NATHAN C: Macrophage deactivation by interleukin 10. *J Exp Med* 174:1548–1555, 1991
193. HOWARD M, O'GARRA A: Biological properties of interleukin 10. *Immunol Today* 13:198–200, 1992
194. JOYCE DA, GIBBONS DP, GREEN P, STEER JH, FELDMANN M, BRENNAN FM: Two inhibitors of pro-inflammatory cytokine release, interleukin-10 and interleukin-4, have contrasting effects on the release of soluble p75 tumor necrosis factor receptor by cultured monocytes. *Eur J Immunol* 24:2699–2705, 1994
195. KITCHING AR, TIPPING PG, HUANG XR, MUTCH DA, HOLDSWORTH SR: Interleukin-4 and interleukin-10 attenuate established crescentic glomerulonephritis in mice. *Kidney Int* 52:52–59, 1997
196. CHADBAN SJ, TESCH GH, FOTI R, ATKINS RC, NIKOLIC-PATERSON DJ: Interleukin-10 is a mesangial cell growth factor in vitro and in vivo. *Lab Invest* 76:619–627, 1997
197. FOUQUERAY B, BOUTARD V, PHILIPPE C, KORNREICH A, MARCHANT A, PEREZ J, GOLDMAN M, BAUD L: Mesangial cell-derived interleukin-10 modulates mesangial cell response to lipopolysaccharide. *Am J Pathol* 147:176–182, 1995
198. ZURAWSKI G, DE VRIES JE: Interleukin 13, an interleukin 4-like cytokine that acts on monocytes and B cells, but not on T cells. *Immunol Today* 15:19–26, 1994
199. LAKKIS FG, CRUET EN: Cloning of rat interleukin-13 (IL-13) cDNA and analysis of IL-13 gene expression in experimental glomerulonephritis. *Biochem Biophys Res Commun* 197:612–618, 1993
200. TANG WW, QI M, VAN GY, WARINER GP, SAMAL B: Leukemia inhibitory factor ameliorates experimental anti-GBM Ab glomerulonephritis. *Kidney Int* 50:1922–1927, 1996
201. HARTNER A, STERZEL RB, REINDL N, HOCKE GM, FEY GH, GOPPELT-STRUEBE M: Cytokine-induced expression of leukemia inhibitory factor in renal mesangial cells. *Kidney Int* 45:1562–1571, 1994
202. LIANOS EA: Eicosanoids in immune-mediated renal injury. *Semin Nephrol* 12:441–453, 1992
203. O'MEARA YM, BRADY HR: Lipoxins, leukocyte recruitment and the resolution phase of acute glomerulonephritis. *Kidney Int* 51(Suppl 58):S56–S61, 1997
204. DIAZ-PEREZ JL, GOLDBYNE ME, WINKELMANN RK: Prostaglandins and chemotaxis: Enhancement of polymorphonuclear leukocyte chemotaxis by prostaglandin F $_{2\alpha}$ . *J Invest Dermatol* 66:149–152, 1976
205. GOODWIN JS, CEUPPENS J: Regulation of the immune response by prostaglandins. *J Clin Immunol* 3:295–315, 1983
206. WEBB DR, OSHEROFF PL: Antigen stimulation of prostaglandin synthesis and control of immune responses. *Proc Natl Acad Sci USA* 73:1300–1304, 1976
207. BRUNDA MJ, HERBERMAN RB, HOLDEN HT: Inhibition of murine natural killer cell activity by prostaglandins. *J Immunol* 124:2682–2687, 1980
208. FANTONE JC, MARASCO WA, ELGAS LJ, WARD PA: Anti-inflammatory effects of prostaglandin E $_1$ : In vivo modulation of the formyl peptide chemotactic receptor on the rat neutrophil. *J Immunol* 130:1495–1497, 1983
209. KAMBAYASHI T, ALEXANDER HR, FONG M, STRASSMANN G: Potential involvement of IL-10 in suppressing tumor-associated macrophages: Colon-26-derived prostaglandin E $_2$  inhibits TNF- $\alpha$  release via a mechanism involving IL-10. *J Immunol* 154:3383–3390, 1995
210. ARMSTRONG RA: Investigation of the inhibitory effects of PGE $_2$  and selective EP agonists on chemotaxis of human neutrophils. *Br J Pharmacol* 116:2903–2908, 1995
211. KUNKEL SL, CHENSUE SW, PHAN SH: Prostaglandins as endogenous mediators of interleukin 1 production. *J Immunol* 136:186–192, 1986
212. MONCADA S, VANE JR: The role of prostacycline in vascular tissue. *Fed Proc* 38:66–71, 1979
213. MENE P, DUNN MJ: Prostaglandins and rat glomerular mesangial cell proliferation. *Kidney Int* 37:1256–1262, 1990
214. ZAHNER G, DISSER M, THAISS F, WOLF G, SCHOEPPE W, STAHL RA: The effect of prostaglandin E $_2$  on mRNA expression and



- secretion of collagens I, III, and IV and fibronectin in cultured rat mesangial cells. *J Am Soc Nephrol* 4:1778–1785, 1994
215. NITTA K, SIMONSON MS, DUNN MJ: The regulation and role of prostaglandin biosynthesis in cultured bovine glomerular endothelial cells. *J Am Soc Nephrol* 2:156–163, 1991
  216. PETRULIS AS, AIKAWA M, DUNN MJ: Prostaglandin and thromboxane synthesis by rat glomerular epithelial cells. *Kidney Int* 20:469–474, 1981
  217. SCHARSCHMIDT LA, DUNN MJ: Prostaglandin synthesis by rat glomerular mesangial cells in culture: Effects of angiotensin II and arginine vasopressin. *J Clin Invest* 71:1756–1764, 1983
  218. LIANOS EA, BRESNAHAN BA, PAN C: Mesangial cell immune injury: Synthesis, origin, and role of eicosanoids. *J Clin Invest* 88:623–631, 1991
  219. FLOEGE J, TOPLEY N, WESSEL K, KAEVER V, RADEKE H, HOPPE J, KISHIMOTO T, RESCH K: Monokines and platelet-derived growth factor modulate prostanoid production in growth arrested, human mesangial cells. *Kidney Int* 37:859–869, 1990
  220. SCHARSCHMIDT LA, DUNN MJ: Prostaglandin synthesis by rat glomerular mesangial cells in culture: Effects of angiotensin II and arginine vasopressin. *J Clin Invest* 71:1756–1764, 1983
  221. HUGHES AK, PADILLA E, KUTCHERA WA, MICHAEL JR, KOHAN DE: Endothelin-1 induction of cyclooxygenase-2 expression in rat mesangial cells. *Kidney Int* 47:53–61, 1995
  222. ALBRIGHTSON CR, NAMBI P, ZABKO-POTAPOVICH B, DYTOK G, GROOM T: Effect of thrombin on proliferation, contraction and prostaglandin production of rat glomerular mesangial cells in culture. *J Pharmacol Exp Ther* 263:404–412, 1992
  223. ADLER S, STAHL RA, BAKER PJ, CHEN YP, PRITZL PM, COUSER WG: Biphasic effect of oxygen radicals on prostaglandin production by rat mesangial cells. *Am J Physiol* 252:F743–F749, 1987
  224. TETSUKA T, DAPHNA-IKEN D, MILLER BW, GUAN Z, BAIER LD, MORRISON AR: Nitric oxide amplifies interleukin 1-induced cyclooxygenase-2 expression in rat mesangial cells. *J Clin Invest* 97:2051–2056, 1996
  225. NEUWIRTH R, SINGHAL P, DIAMOND B, HAYS RM, LOBMEYER L, CLAY K, SCHLÖNDORFF D: Evidence for immunoglobulin Fc receptor-mediated prostaglandin and platelet-activating factor formation by cultured rat mesangial cells. *J Clin Invest* 82:936–944, 1988
  226. LIEBERTHAL W, LEVINE L: Stimulation of prostaglandin production in rat glomerular epithelial cells by antidiuretic hormone. *Kidney Int* 25:766–770, 1984
  227. LIANOS EA, ANDRES GA, DUNN MJ: Glomerular prostaglandin and thromboxane synthesis in rat nephrotoxic serum nephritis: Effects on renal hemodynamics. *J Clin Invest* 72:1439–1448, 1983
  228. STAHL RA, ADLER S, BAKER PJ, CHEN YP, PRITZL PM, COUSER WG: Enhanced glomerular prostaglandin formation in experimental membranous nephropathy. *Kidney Int* 31:1126–1131, 1987
  229. McLEISH KR, GOHARA AF, STELZER GT, WALLACE JH: Treatment of murine immune complex glomerulonephritis with prostaglandin E<sub>2</sub>: Dose-response of immune complex deposition, antibody synthesis, and glomerular damage. *Clin Immunol Immunopathol* 26:18–23, 1983
  230. McLEISH KR, STELZER GT, EADES DS, COHEN R, WALLACE JH: Serial changes in humoral and cellular immunity induced by prostaglandin E<sub>2</sub> treatment of murine immune complex glomerulonephritis. *J Lab Clin Med* 106:517–523, 1985
  231. POELSTRA K, BROUWER E, BALLER JF, HARDONK MJ, BAKKER WW: Attenuation of anti-Thy1 glomerulonephritis in the rat by anti-inflammatory platelet-inhibiting agents. *Am J Pathol* 142:441–450, 1993
  232. LEE TH, HORTON CE, KYAN-AUNG U, HASKARD D, CREA AE, SPUR BW: Lipoxin A<sub>4</sub> and lipoxin B<sub>4</sub> inhibit chemotactic responses of human neutrophils stimulated by leukotriene B<sub>4</sub> and N-formyl-L-methionyl-L-leucyl-L-phenylalanine. *Clin Sci* 77:195–203, 1989
  233. PAPAYIANNI A, SERHAN CN, BRADY HR: Lipoxin A<sub>4</sub> and B<sub>4</sub> inhibit leukotriene-stimulated interactions of human neutrophils and endothelial cells. *J Immunol* 156:2264–2272, 1996
  234. PAPAYIANNI A, SERHAN CN, PHILLIPS ML, RENNKE HG, BRADY HR: Transcellular biosynthesis of lipoxin A<sub>4</sub> during adhesion of platelets and neutrophils in experimental immune complex glomerulonephritis. *Kidney Int* 47:1295–1302, 1995
  235. MAYADAS TN, MENDRICK DL, BRADY HR, TANG T, PAPAYIANNI A, ASSMANN KJ, WAGNER DD, HYNES RO, COTRAN RS: Acute passive anti-glomerular basement membrane nephritis in P-selectin-deficient mice. *Kidney Int* 49:1342–1349, 1996
  236. GARRICK R, SHEN SY, OGUNC S, WONG PY: Transformation of leukotriene A<sub>4</sub> to lipoxins by rat kidney mesangial cell. *Biochem Biophys Res Commun* 162:626–633, 1989
  237. GARRICK R, GOODMAN A, SHEN ST, OGUNC S, WONG PY: Regulation of lipoxins and leukotriene B<sub>4</sub> production in rat mesangial cells. *Adv Prostaglandin Thromb Leuk Res* 21B:701–706, 1991
  238. KATOH T, LAKKIS FG, MAKITA N, BADR KF: Co-regulated expression of glomerular 12/15-lipoxygenase and interleukin-4 mRNAs in rat nephrotoxic nephritis. *Kidney Int* 46:341–349, 1994
  239. BRADY HR, PERSSON U, BALLERMANN BJ, BRENNER BM, SERHAN CN: Leukotrienes stimulate neutrophil adhesion to mesangial cells: Modulation with lipoxins. *Am J Physiol* 259:F809–F815, 1990
  240. KELEFIOTIS D, BRESNAHAN BA, STRATIDAKIS I, LIANOS EA: Eicosanoid-induced growth and signaling events in rat glomerular mesangial cells. *Prostaglandins* 49:269–283, 1995
  241. BADR KF, DEBOER DK, SCHWARTZBERG M, SERHAN CN: Lipoxin A<sub>4</sub> antagonizes cellular and in vivo actions of leukotriene D<sub>4</sub> in rat glomerular mesangial cells: Evidence for competition at a common receptor. *Proc Natl Acad Sci USA* 86:3438–3442, 1989
  242. VOLL RE, HERRMANN M, ROTH EA, STACH C, KALDEN JR, GIRONKONTAITE I: Immunosuppressive effects of apoptotic cells. *Nature* 390:350–351, 1997
  243. FADOK VA, BRATTON DL, KONOWAL A, FREED PW, WESTCOTT JY, HENSON PM: Macrophages that have ingested apoptotic cells in vitro inhibit proinflammatory cytokine production through autocrine/paracrine mechanisms involving TGF- $\beta$ , PGE<sub>2</sub>, and PAF. *J Clin Invest* 101:890–898, 1998
  244. HARRISON DJ: Cell death in the diseased glomerulus. *Histopathology* 12:679–683, 1988
  245. BAKER AJ, MOONEY A, HUGHES J, LOMBARDI D, JOHNSON RJ, SAVILL J: Mesangial cell apoptosis: The major mechanism for resolution of glomerular hypercellularity in experimental mesangial proliferative nephritis. *J Clin Invest* 94:2105–2116, 1994
  246. TAKEMURA T, MURAKAMI K, MIYAZATO H, YAGI K, YOSHIOKA K: Expression of Fas and Bcl-2 in human glomerulonephritis. *Kidney Int* 48:1886–1892, 1995
  247. SHIMIZU A, MASUDA Y, KITAMURA H, ISHIZAKI M, SUGISAKI Y, YAMANAKA N: Apoptosis in progressive crescentic glomerulonephritis. *Lab Invest* 74:941–951, 1996
  248. SUGIYAMA H, KASHIHARA N, MAKINO H, YAMASAKI Y, OTA Z: Apoptosis in glomerular sclerosis. *Kidney Int* 49:103–111, 1996
  249. KITAMURA M, MITARAI T, MARUYAMA N, NAGASAWA R, YOSHIDA H, SAKAI O: Mesangial cell behavior in a three-dimensional extracellular matrix. *Kidney Int* 40:653–661, 1991
  250. YAOITA E: Behavior of rat mesangial cells cultured within extracellular matrix. *Lab Invest* 61:410–418, 1989
  251. KITAMURA M, ISHIKAWA Y: Three-dimensional matrix primes mesangial cells to down-regulation of  $\alpha$ -smooth muscle actin via deactivation of CArG box elements. *Kidney Int* 53:690–697, 1998
  252. KITAMURA M, MITARAI T, NAGASAWA R, MARUYAMA N: Differentiated phenotype of glomerular mesangial cells in nodular culture. *Am J Physiol* 270:F614–F622, 1996
  253. TOYABE S, IWANAGA T: An ultrastructural study of proliferative nephritis induced experimentally by a monoclonal antibody against mesangial cells. *Virchows Arch B Cell Pathol* 61:397–407, 1992
  254. STEINMANN-NIGGLI K, ZISWILER R, KUNG M, MARTI H-P: Inhibition of matrix metalloproteinases attenuates anti-Thy1.1 nephritis. *J Am Soc Nephrol* 9:397–407, 1998
  255. MARX M, DANIEL TO, KASHGARIAN M, MADRI JA: Spatial organization of the extracellular matrix modulates the expression of PDGF-receptor subunits in mesangial cells. *Kidney Int* 43:1027–1041, 1993
  256. JOHNSON RJ: Platelets in inflammatory glomerular injury. *Semin Nephrol* 11:276–284, 1991
  257. DeBAULT LE, ESMON NL, OLSON JR, ESMON CT: Distribution of the thrombomodulin antigen in the rabbit vasculature. *Lab Invest* 54:172–178, 1986
  258. MIZUTANI M, YUZAWA Y, MARUYAMA I, SAKAMOTO N, MATSUO S:

- Glomerular localization of thrombomodulin in human glomerulonephritis. *Lab Invest* 69:193–202, 1993
259. HE CJ, KANFER A: Quantification and modulation of thrombomodulin activity in isolated rat and human glomeruli. *Kidney Int* 41:1170–1174, 1992
  260. PRUNA A, PEYRI N, BERARD M, BOFFA MC: Thrombomodulin is synthesized by human mesangial cells. *Kidney Int* 51:687–693, 1997
  261. GONDA Y, HIRATA S, SAITOH K, AOKI Y, MOHRI M, GOMI K, SUGIHARA T, KIIYOTA T, YAMAMOTO S, ISHIDA T: Antithrombotic effect of recombinant human soluble thrombomodulin on endotoxin-induced disseminated intravascular coagulation in rats. *Thromb Res* 71:325–335, 1993
  262. BREGENGARD C, NORDFANG O, WILDGOOSE P, SVENDSEN O, HEDNER U, DINESS V: The effect of two-domain tissue factor pathway inhibitor on endotoxin-induced disseminated intravascular coagulation in rabbits. *Blood Coagul Fibrinolysis* 4:699–706, 1993
  263. ERLICH JH, APOSTOLOPOULOS J, WUN TC, KRETZMER KK, HOLDSWORTH SR, TIPPING PG: Renal expression of tissue factor pathway inhibitor and evidence for a role in crescentic glomerulonephritis in rabbits. *J Clin Invest* 98:325–335, 1996
  264. YAMABE H, OSAWA H, INUMA H, KAIZUKA M, TAMURA N, TSUNODA S, FUJITA Y, SHIRATO K, ONODERA K: Tissue factor pathway inhibitor production by human mesangial cells in culture. *Thromb Haemost* 76:215–219, 1996
  265. MALLIAROS J, HOLDSWORTH SR, WOJTA J, ERLICH J, TIPPING PG: Glomerular fibrinolytic activity in anti-GBM glomerulonephritis in rabbits. *Kidney Int* 44:557–564, 1993
  266. ZOJA C, CORNA D, MACCONI D, ZILIO P, BERTANI T, REMUZZI G: Tissue plasminogen activator therapy of rabbit nephrotoxic nephritis. *Lab Invest* 62:34–40, 1990
  267. ANGLES-CANO E, RONDEAU E, DELARUE F, HAGEGE J, SULTAN Y, SRAER JD: Identification and cellular localization of plasminogen activators from human glomeruli. *Thromb Haemost* 54:688–692, 1985
  268. AYA N, YOSHIOKA K, MURAKAMI K, HINO S, OKADA K, MATSUO O, MAKI S: Tissue-type plasminogen activator and its inhibitor in human glomerulonephritis. *J Pathol* 166:289–295, 1992
  269. YAMAMOTO K, LOSKUTOFF DJ: The kidneys of mice with autoimmune disease acquire a hypofibrinolytic/procoagulant state that correlates with the development of glomerulonephritis and tissue microthrombosis. *Am J Pathol* 151:725–734, 1997
  270. LACAVE R, RONDEAU E, OCHI S, DELARUE F, SCHLEUNING WD, SRAER JD: Characterization of a plasminogen activator and its inhibitor in human mesangial cells. *Kidney Int* 35:806–811, 1989
  271. IWAMOTO T, NAKASHIMA Y, SUEISHI K: Secretion of plasminogen activator and its inhibitor by glomerular epithelial cells. *Kidney Int* 37:1466–1476, 1990
  272. WILSON HM, HAITES NE, REID FJ, BOOTH NA: Interleukin-1 $\beta$  up-regulates the plasminogen activator/plasmin system in human mesangial cells. *Kidney Int* 49:1097–1104, 1996
  273. KITCHING AR, HOLDSWORTH SR, PLOPLIS VA, PLOW EF, COLLEN D, CARMELIET P, TIPPING PG: Plasminogen and plasminogen activators protect against renal injury in crescentic glomerulonephritis. *J Exp Med* 185:963–968, 1997
  274. STRIKER LJ, PETEN EP, ELLIOT SJ, DOI T, STRIKER GE: Mesangial cell turnover: Effect of heparin and peptide growth factors. *Lab Invest* 64:446–456, 1991
  275. WARDLE EN: Heparins for proliferative nephritides? Short review on an advancing topic. *Nephron* 73:515–519, 1996
  276. MARCUM JA, ATHA DH, FRITZE LM, NAWROTH P, STERN D, ROSENBERG RD: Cloned bovine aortic endothelial cells synthesize anticoagulant active heparan sulfate proteoglycan. *J Biol Chem* 261:7507–7517, 1986
  277. FERRER-LOPEZ P, RENESTO P, PREVOST MC, GOUNON P, CHIGNARD M: Heparin inhibits neutrophil-induced platelet activation via cathepsin G. *J Lab Clin Med* 119:231–239, 1992
  278. YOKOKAWA K, TAHARA H, KOHNO M, MANDAL AK, YANAGISAWA M, TAKEDA T: Heparin regulates endothelin production through endothelium-derived nitric oxide in human endothelial cells. *J Clin Invest* 92:2080–2085, 1993
  279. VAIREL EG, BOUTY-BOYE H, TOULEMONDE F, DOUTREMEPUICH C, MARSH NA, GAFFNEY PJ: Heparin and a low molecular weight fraction enhances thrombolysis and by this pathway exercises a protective effect against thrombosis. *Thromb Res* 30:219–224, 1983
  280. WEILER JM, EDENS RE, LINHARDT RJ, KAPELANSKI DP: Heparin and modified heparin inhibit complement activation in vivo. *J Immunol* 148:3210–3215, 1992
  281. EKRE HP, NAPARSTEK Y, LIDER O, HYDEN P, HAGERMARK O, NILSSON T, VLODAVSKY I, COHEN I: Anti-inflammatory effects of heparin and its derivatives: Inhibition of complement and of lymphocyte migration. *Adv Exp Med Biol* 313:329–340, 1992
  282. CAUGHMAN GB, BOACKLE RJ, VESELY J: A postulated mechanism for heparin's potentiation of C1 inhibitor function. *Mol Immunol* 19:287–295, 1982
  283. KAZATCHKINE MD, FEARON DT, SILBERT JE, AUSTEN KF: Surface-associated heparin inhibits zymosan-induced activation of the human alternative complement pathway by augmenting the regulatory action of the control proteins on particle-bound C3b. *J Exp Med* 150:1202–1215, 1979
  284. KENAGY RD, NIKKARI ST, WELGUS HG, CLOWES AW: Heparin inhibits the induction of three matrix metalloproteinases (stromelysin, 92-kD gelatinase, and collagenase) in primate arterial smooth muscle cells. *J Clin Invest* 93:1987–1993, 1994
  285. WALSH RL, DILLON TJ, SCICCHITANO R, MCLENNAN G: Heparin and heparan sulphate are inhibitors of human leucocyte elastase. *Clin Sci (Colch)* 81:341–346, 1991
  286. LEY K, CERRITO M, ARFORS KE: Sulfated polysaccharides inhibit leukocyte rolling in rabbit mesentery venules. *Am J Physiol* 260:H1667–H1673, 1991
  287. BAZZONI G, BELTRAN-NUNEZ A, MASCELLANI G, BIANCHINI P, DEJANA E, DEL-MASCHIO A: Effect of heparin, dermatan sulfate, and related oligo-derivatives on human polymorphonuclear leukocyte functions. *J Lab Clin Med* 121:268–275, 1993
  288. NELSON RM, CECCONI O, ROBERTS WG, ARUFFO A, LINHARDT RJ, BEVILACQUA MP: Heparin oligosaccharides bind L- and P-selectin and inhibit acute inflammation. *Blood* 82:3253–3258, 1993
  289. GARRED P, MOLLNES TE: Immobilized heparin inhibits the increase in leukocyte surface expression of adhesion molecules. *Artif Organs* 21:293–299, 1997
  290. MATZNER Y, MARX G, DREXLER R, ELDOR A: The inhibitory effect of heparin and related glycosaminoglycans on neutrophil chemotaxis. *Thromb Haemost* 52:134–137, 1984
  291. LAGHI-PASINI F, PASQUI AL, CECCATELLI L, CAPECCHI PL, ORRICO A, DI-PERRI T: Heparin inhibition of polymorphonuclear leukocyte activation in vitro: A possible pharmacological approach to granulocyte-mediated vascular damage. *Thromb Res* 35:527–537, 1984
  292. MAVIER P, PREAUX AM, ROSENBAUM J, ZAFRANI ES, DHUMEAUX D: Toxicity of polymorphonuclear neutrophils against hepatocytes: Protective effect of heparin. *Biochem Pharmacol* 38:1865–1867, 1989
  293. MANASTER J, CHEZAR J, SHURTZ-SWIRSKI R, SHAPIRO G, TENDLER Y, KRISTAL B, SHASHA SM, SELA S: Heparin induces apoptosis in human peripheral blood neutrophils. *Br J Haematol* 94:48–52, 1996
  294. LEUNG L, SAIGO K, GRANT D: Heparin binds to human monocytes and modulates their procoagulant activities and secretory phenotypes: Effects of histidine-rich glycoprotein. *Blood* 73:177–184, 1989
  295. CASTELLOT JJ JR, HOOVER RL, HARPER PA, KARNOVSKY MJ: Heparin and glomerular epithelial cell-secreted heparin-like species inhibit mesangial-cell proliferation. *Am J Pathol* 120:427–435, 1985
  296. CASTELLOT JJ JR, HOOVER RL, KARNOVSKY MJ: Glomerular endothelial cells secrete a heparinlike inhibitor and a peptide stimulator of mesangial cell proliferation. *Am J Pathol* 125:493–500, 1986
  297. GROGGER GC, MARINIDES GN, HOVINGH P, HAMMOND E, LINKER A: Inhibition of rat mesangial cell growth by heparan sulfate. *Am J Physiol* 258:F259–F265, 1990
  298. PERSON JM, LOVETT DH, RAUGI GJ: Modulation of mesangial cell migration by extracellular matrix components: Inhibition by heparinlike glycosaminoglycans. *Am J Pathol* 133:609–614, 1988
  299. MIRALEM T, WANG A, WHITESIDE CI, TEMPLETON DM: Heparin

- inhibits mitogen-activated protein kinase-dependent and -independent *c-fos* induction in mesangial cells. *J Biol Chem* 271:17100–17106, 1996
300. HOFER G, GRIMMER C, SUKHATME VP, STERZEL RB, RUPPRECHT HD: Transcription factor Egr-1 regulates glomerular mesangial cell proliferation. *J Biol Chem* 271:28306–29310, 1996
  301. CAENAZZO C, GARBISA S, CEOL M, BAGGIO B, BORSATTI A, MARCHI E, GAMBARO G: Heparin modulates proliferation and proteoglycan biosynthesis in murine mesangial cells: Molecular clues for its activity in nephropathy. *Nephrol Dial Transplant* 10:175–184, 1995
  302. ISHIKAWA Y, KITAMURA M: Spontaneous apoptosis of podocytes in isolated glomeruli. *Kidney Int* 54:2008–2013, 1998
  303. ADLER S: Inhibition of rat glomerular visceral epithelial cell growth by heparin. *Am J Physiol* 255:F781–F786, 1988
  304. ADLER S: Heparin alters epidermal growth factor metabolism in cultured rat glomerular epithelial cells. *Am J Pathol* 139:169–175, 1991
  305. KARLSSON K, MARKLUND SL: Heparin-induced release of extracellular superoxide dismutase to human blood plasma. *Biochem J* 242:55–59, 1987
  306. HIEBERT LM, LIU JM: Heparin protects cultured arterial endothelial cells from damage by toxic oxygen metabolites. *Atherosclerosis* 83:47–51, 1990
  307. KANWAR YS, ROSENZWEIG LJ, JAKUBOWSKI ML: Distribution of de novo synthesized sulfated glycosaminoglycans in the glomerular basement membrane and mesangial matrix. *Lab Invest* 49:216–225, 1983
  308. GROFFEN AJ, HOP FW, TRYGGVASON K, DIJKMAN H, ASSMANN KJ, VEERKAMP JH, MONNENS LA, VAN DEN HEUVEL LP: Evidence for the existence of multiple heparan sulfate proteoglycans in the human glomerular basement membrane and mesangial matrix. *Eur J Biochem* 247:175–182, 1997
  309. RUPPRECHT HD, SCHOCKLMANN HO, STERZEL RB: Cell-matrix interaction in the glomerular mesangium. *Kidney Int* 49:1575–1582, 1996
  310. THOMAS GJ, MASON RM, DAVIES M: Characterization of proteoglycans synthesized by human adult glomerular mesangial cells in culture. *Biochem J* 277:81–88, 1991
  311. VAN DET NF, VAN DEN BORN J, TAMSMA JT, VERHAGEN NA, VAN DEN HEUVEL LP, BERDEN JH, BRUIJN JA, DAHA MR, VAN DER WOUDE FJ: Proteoglycan production by human glomerular visceral epithelial cells and mesangial cells in vitro. *Biochem J* 307:759–768, 1995
  312. KOLM V, SAUER U, OLGEMOOLLER B, SCHLEICHER ED: High glucose-induced TGF- $\beta$ 1 regulates mesangial production of heparan sulfate proteoglycan. *Am J Physiol* 270:F812–F821, 1996
  313. VAN DET NF, TAMSMA JT, VAN DEN BORN J, VERHAGEN NA, VAN DEN HEUVEL LP, LOWIK CW, BERDEN JH, BRUIJN JA, DAHA MR, VAN DER WOUDE FJ: Differential effects of angiotensin II and transforming growth factor- $\beta$  on the production of heparan sulfate proteoglycan by mesangial cells in vitro. *J Am Soc Nephrol* 7:1015–1023, 1996
  314. RAIJ L, BAYLIS C: Glomerular actions of nitric oxide. *Kidney Int* 48:20–32, 1995
  315. PRYOR WA, SQUADRITO GL: The chemistry of peroxynitrite: A product from the reaction of nitric oxide with superoxide. *Am J Physiol* 268:L699–L722, 1995
  316. SHULTZ PJ, TAYEH MA, MARLETTA MA, RAIJ L: Synthesis and action of nitric oxide in rat glomerular mesangial cells. *Am J Physiol* 261:F600–F606, 1991
  317. PFEILSCHIFTER J, ROB P, MULSCH A, FANDREY J, VOSBECK K, BUSSE R: Interleukin 1- $\beta$  and tumour necrosis factor- $\alpha$  induce a macrophage-type of nitric oxide synthase in rat renal mesangial cells. *Eur J Biochem* 203:251–255, 1992
  318. SANDAU K, PFEILSCHIFTER J, BRUNE B: The balance between nitric oxide and superoxide determines apoptotic and necrotic death of rat mesangial cells. *J Immunol* 158:4938–4946, 1997
  319. KUBES P, SUZUKI M, GRANGER DN: Nitric oxide: An endogenous modulator of leukocyte adhesion. *Proc Natl Acad Sci USA* 88:4651–4655, 1991
  320. CLANCY RM, LESZCZYNSKA-PIZIAK J, ABRAMSON SB: Nitric oxide, an endothelial cell relaxation factor, inhibits neutrophil superoxide anion production via a direct action on the NADPH oxidase. *J Clin Invest* 90:1116–1121, 1992
  321. SU WY, DAY BJ, KANG BH, CRAPO JD, HUANG YC, CHANG LY: Lung epithelial cell-released nitric oxide protects against PMN-mediated cell injury. *Am J Physiol* 271:L581–L586, 1996
  322. RADOMSKI MW, PALMER RM, MONCADA S: Comparative pharmacology of endothelium-derived relaxing factor, nitric oxide and prostacyclin in platelets. *Br J Pharmacol* 92:181–187, 1987
  323. RADOMSKI MW, PALMER RM, MONCADA S: Endogenous nitric oxide inhibits human platelet adhesion to vascular endothelium. *Lancet* 2:1057–1058, 1987
  324. SHULTZ PJ, RAIJ L: Endogenously synthesized nitric oxide prevents endotoxin-induced glomerular thrombosis. *J Clin Invest* 90:1718–1725, 1992
  325. WESTBERG G, SHULTZ PJ, RAIJ L: Exogenous nitric oxide prevents endotoxin-induced glomerular thrombosis in rats. *Kidney Int* 46:711–716, 1994
  326. MUHL H, SANDAU K, BRUNE B, BRINER VA, PFEILSCHIFTER J: Nitric oxide donors induce apoptosis in glomerular mesangial cells, epithelial cells and endothelial cells. *Eur J Pharmacol* 317:137–149, 1996
  327. GARG UC, HASSID A: Inhibition of rat mesangial cell mitogenesis by nitric oxide-generating vasodilators. *Am J Physiol* 257:F60–F66, 1989
  328. CRAVEN PA, STUDER RK, FELDER J, PHILLIPS S, DERUBERTIS FR: Nitric oxide inhibition of transforming growth factor- $\beta$  and collagen synthesis in mesangial cells. *Diabetes* 46:671–681, 1997
  329. IKEDA M, IKEDA U, TAKAHASHI M, SHIMADA K, MINOTA S, KANO S: Nitric oxide inhibits intracellular adhesion molecule-1 expression in rat mesangial cells. *J Am Soc Nephrol* 7:2213–2218, 1996
  330. ARMSTEAD VE, MINCHENKO AG, SCHUHL RA, HAYWARD R, NOS-SULI TO, LEFER AM: Regulation of P-selectin expression in human endothelial cells by nitric oxide. *Am J Physiol* 273:H740–H746, 1997
  331. HABERSTROH U, STILO K, POCKOCK J, WOLF G, HELMCHEN U, WENZEL U, ZAHNER G, STAHL RAK, THAISS F: L-Arginine suppresses lipopolysaccharide-induced expression of RANTES in glomeruli. *J Am Soc Nephrol* 9:203–210, 1998
  332. WELCH WJ: Mammalian stress response: Cell physiology, structure/function of stress proteins, and implications for medicine and disease. *Physiol Rev* 72:1063–1081, 1992
  333. YOKOO T, KITAMURA M: Heat shock protein in the kidney. *Exp Nephrol* 5:439–444, 1997
  334. HARTL FU: Molecular chaperones in cellular protein folding. *Nature* 381:571–579, 1996
  335. CRAIG EA, WEISSMAN JS, HORWICH AL: Heat shock proteins and molecular chaperones: Mediators of protein conformation and turnover in the cells. *Cell* 78:365–372, 1994
  336. KOMATSUDA A, WAKUI H, IMAI H, NAKAMOTO Y, MIURA AB, ITOH H, TASHIMA Y: Renal localization of the constitutive 73-kDa heat-shock protein in normal and PAN rats. *Kidney Int* 41:1204–1212, 1992
  337. YOKOO T, KITAMURA M: IL-1 $\beta$  depressed expression of the 70-kilodalton heat shock protein and sensitizes glomerular cells to oxidant-initiated apoptosis. *J Immunol* 159:2886–2892, 1997
  338. GABAI VL, MERIIN AB, MOSSER DD, CARON AW, RITS S, SHIFRIN VI, SHERMAN MY: Hsp70 prevents activation of stress kinases: A novel pathway of cellular thermotolerance. *J Biol Chem* 272:18033–18037, 1997
  339. ISHIKAWA Y, YOKOO T, KITAMURA M: c-Jun/AP-1, but not NF- $\kappa$ B, is a mediator for oxidant-initiated apoptosis in glomerular mesangial cells. *Biochem Biophys Res Commun* 240:496–501, 1997
  340. WAKUI H, ITOH H, TASHIMA Y, KOBAYASHI R, NAKAMOTO Y, MIURA AB: Purification of a mature form of 60 kDa heat-shock protein (chaperonin homolog) from porcine kidney and its partial amino acid sequence. *Int J Biochem* 24:1507–1510, 1992
  341. MATSUBARA O, KASUGA T, MARUMO F, ITOH H, TASHIMA Y: Localization of 90-kDa heat shock protein in the kidney. *Kidney Int* 38:830–834, 1990
  342. MOLITERNO R, WOAN M, BENTLEJEWSKI C: Heat shock protein-induced T lymphocyte propagation from endomyocardial biopsies in heart transplantation. *J Heart Lung Transplant* 14:329–337, 1995



343. GALEA-LAURI J, RICHARDSON AJ, LATCHMAN DS, KATZ DR: Increased heat shock protein 90 (hsp90) expression leads to increased apoptosis in the monoblastoid cell line U937 following induction with TNF- $\alpha$  and cycloheximide: A possible role in immunology. *J Immunol* 157:4109–4118, 1996
344. KAHN W, MCGUIRT JP, SENS MA, SENS DA, TODD JH: Expression of heat shock protein 27 in developing and adult human kidney. *Toxicol Lett* 84:69–79, 1996
345. SMOYER WE, GUPTA A, MUNDEL P, BALLEW JD, WELSH MJ: Altered expression of glomerular heat shock protein 27 in experimental nephrotic syndrome. *J Clin Invest* 97:2697–2704, 1996
346. MAINES MD: The heme oxygenase system: A regulator of second messenger gases. *Annu Rev Pharmacol Toxicol* 37:517–554, 1997
347. APPELGATE LA, LUSCHER P, TYRRELL RM: Induction of heme oxygenase: A general response to oxidant stress in cultured mammalian cells. *Cancer Res* 51:974–978, 1991
348. STOCKER R: Induction of haem oxygenase as a defense against oxidative stress. *Free Radic Res Commun* 9:101–112, 1990
349. LEE PJ, ALAM J, WIEGAND GW, CHOI AM: Overexpression of heme oxygenase-1 in human pulmonary epithelial cells results in cell growth arrest and increased resistance to hyperoxia. *Proc Natl Acad Sci USA* 93:10393–10398, 1996
350. WILLIS D, MOORE AR, FREDERICK R, WILLOUGHBY DA: Heme oxygenase: A novel target for the modulation of the inflammatory response. *Nat Med* 2:87–90, 1996
351. LANIADO-SCHWARTZMAN M, ABRAHAM NG, CONNERS M, DUNN MW, LEVERE RD, KAPPAS A: Heme oxygenase induction with attenuation of experimentally induced corneal inflammation. *Biochem Pharmacol* 53:1069–1075, 1997
352. TETSUKA T, DAPHNA-IKEN D, SRIVASTAVA SK, MORRISON AR: Regulation of heme oxygenase mRNA in mesangial cells: Prostaglandin E2 negatively modulates interleukin-1-induced heme oxygenase-1 mRNA. *Biochem Biophys Res Commun* 212:617–623, 1995
353. SCHENA FP, GESUALDO L, GRANDALIANO G, MONTINARO V: Progression of renal damage in human glomerulonephritides: Is there sleight of hand in winning the game? *Kidney Int* 52:1439–1457, 1997
354. YOSHIOKA T, ICHIKAWA I, FOGO A: Reactive oxygen metabolites cause massive, reversible proteinuria and glomerular sieving defect without apparent ultrastructural abnormality. *J Am Soc Nephrol* 2:902–912, 1991
355. NAKAMURA H, NAKAMURA K, YODOI J: Redox regulation of cellular activation. *Annu Rev Immunol* 15:351–369, 1997
356. SUZUKI YJ, FORMAN HJ, SEVANIAN A: Oxidants as stimulators of signal transduction. *Free Radic Biol Med* 22:269–285, 1997
357. MAYER TN, GLOY J, HUG MJ, GREGER R, SCHOLLMAYER P, PAVENSTADT H: Hydrogen peroxide increases the intracellular calcium activity in rat mesangial cells in primary culture. *Kidney Int* 49:388–395, 1996
358. YOKOO T, KITAMURA M: Unexpected protection of glomerular mesangial cells from oxidant-triggered apoptosis by bioflavonoid quercetin. *Am J Physiol* 273:F206–F212, 1997
359. ANGEL P, KARIN M: The role of Jun, Fos and the AP-1 complex in cell-proliferation and transformation. *Biochim Biophys Acta* 1072:129–157, 1991
360. BAEUERLE PA, HENKEL T: Function and activation of NF- $\kappa$ B in the immune system. *Annu Rev Immunol* 12:147–179, 1994
361. SATRIANO JA, SHULDINER M, HORA K, XING Y, SHAN Z, SCHLÖNDORFF D: Oxygen radicals as second messengers for expression of the monocyte chemoattractant protein, JE/MCP-1, and the monocyte colony-stimulating factor, CSF-1, in response to tumor necrosis factor- $\alpha$  and immunoglobulin G: Evidence for involvement of reduced nicotinamide adenine dinucleotide phosphate (NADPH)-dependent oxidase. *J Clin Invest* 92:1564–1571, 1993
362. LO SK, JANAKIDEVI K, LAI L, MALIK AB: Hydrogen peroxide-induced increase in endothelial adhesiveness is dependent on ICAM-1 activation. *Am J Physiol* 264:L406–L412, 1993
363. ICHIKAWA I, KIYAMA S, YOSHIOKA T: Renal antioxidant enzymes: Their regulation and function. *Kidney Int* 45:1–9, 1994
364. BANNISTER JV, BANNISTER WH, ROTILIO G: Aspects of the structure, function, and applications of superoxide dismutase. *CRC Crit Rev Biochem* 22:111–180, 1987
365. PRZEDBORSKI S, KOSTIC V, JACKSON-LEWIS V, NAINI AB, SIMONETTI S, FAHN S, CARLSON E, EPSTEIN CJ, CADET JL: Transgenic mice with increased Cu/Zn-superoxide dismutase activity are resistant to N-methyl-4-phenyl-1,2,3,6-tetrahydropyridine-induced neurotoxicity. *J Neurosci* 12:1658–1667, 1992
366. WISPE JR, WARNER BB, CLARK JC, DEY CR, NEUMAN J, GLASSER SW, CRAPO JD, CHANG L-Y, WHITSETT JA: Human Mn-superoxide dismutase in pulmonary epithelial cells of transgenic mice confers protection from oxygen injury. *J Biol Chem* 267:23937–23941, 1992
367. YOSHIOKA T, BILLS T, MOORE-JARRETT T, GREENE HL, BURR IM, ICHIKAWA I: Role of intrinsic antioxidant enzymes in renal oxidant injury. *Kidney Int* 38:282–288, 1990
368. YOSHIOKA T, KAWAMURA T, MEYRICK BO, BECKMAN JK, HOOVER RL, YOSHIDA H, ICHIKAWA I: Induction of manganese superoxide dismutase by glucocorticoids in glomerular cells. *Kidney Int* 45:211–219, 1994
369. GWINNER W, TISHER CC, NICK HS: Regulation of manganese superoxide dismutase in glomerular epithelial cells: Mechanisms for interleukin-1 induction. *Kidney Int* 48:354–362, 1995
370. DIAMOND JR, BONVENTRE JV, KARNOVSKY MJ: A role for oxygen free radicals in aminonucleoside nephrosis. *Kidney Int* 29:478–483, 1986
371. BEAMAN M, BIRTWISTLE R, HOWIE AJ, MICHAEL J, ADU D: The role of superoxide anion and hydrogen peroxide in glomerular injury induced by puromycin aminonucleoside in rats. *Clin Sci* 73:329–332, 1987
372. YOSHIOKA T, HOMMA T, MEYRICK B, TAKEDA M, MOORE-JARRETT T, KON V, ICHIKAWA I: Oxidants induce transcriptional activation of manganese superoxide dismutase in glomerular cells. *Kidney Int* 46:405–413, 1994
373. STEPHANZ GB, GWINNER W, CANNON JK, TISHER CC, NICK HS: Heat-aggregated IgG and interleukin-1 $\beta$  stimulate manganese superoxide dismutase in mesangial cells. *Exp Nephrol* 4:151–158, 1996
374. ADACHI T, FUKUTA M, ITO Y, HIRANO K, SUGIURA M, SUGIURA K: Effect of superoxide dismutase on glomerular nephritis. *Biochem Pharmacol* 35:341–345, 1986
375. LINDAU-SHEPARD BA, SHAFFER JB: Expression of human catalase in acatalasemic murine SV-B2 cells confers protection from oxidative damage. *Free Radic Biol Med* 15:581–588, 1993
376. LIEBMANN J, FISHER J, LIPSCHULTZ C, KUNO R, KAUFMAN DC: Enhanced glutathione peroxidase expression protects cells from hydroperoxides but not from radiation or doxorubicin. *Cancer Res* 55:4465–4470, 1995
377. BALIGA R, BALIGA M, SHAH SV: Effect of selenium deficient diet in experimental glomerular disease. *Am J Physiol* 263:F56–F61, 1992
378. NATH KA, SALAHUDEEN AK: Induction of renal growth and injury in the intact rat kidney by dietary deficiency of antioxidants. *J Clin Invest* 86:1179–1192, 1990
379. HUNTER T: Protein kinases and phosphatases: The Yin and Yang of protein phosphorylation and signaling. *Cell* 80:225–236, 1995
380. WHITMARSH AJ, DAVIS RJ: Transcription factor AP-1 regulation by mitogen-activated protein kinase signal transduction pathways. *J Mol Med* 74:589–607, 1996
381. BOKEMEYER D, SOROKIN A, DUNN MJ: Differential regulation of the dual-specificity protein-tyrosine phosphatases CL100, B23, and PAC1 in mesangial cells. *J Am Soc Nephrol* 8:40–50, 1997
382. WENZEL UO, FOUQUERAY B, BISWAS P, GRANDALIANO G, CHOUDHURY GG, ABBODD HE: Activation of mesangial cells by the phosphatase inhibitor vanadate: Potential implications for diabetic nephropathy. *J Clin Invest* 95:1244–1252, 1995
383. FENG L, XIA Y, SEIFFERT D, WILSON CB: Oxidative stress-inducible protein tyrosine phosphatase in glomerulonephritis. *Kidney Int* 48:1920–1928, 1995
384. CHENG J, WU K, ARMANINI M, O'Rourke N, DOWBENKO D, LASKY LA: A novel protein-tyrosine phosphatase related to the homotypically adhering kappa and mu receptors. *J Biol Chem* 272:7264–7277, 1997
385. TAGAWA M, SHIRASAWA T, YAHAGI Y, TOMODA T, KUROYANAGI H, FUJIMURA S, SAKIYAMA S, MARUYAMA N: Identification of a receptor-type protein-tyrosine phosphatase expressed in postmi-



- totot maturing neurons: Its structure and expression in the central nervous system. *Biochem J* 321:865–871, 1997
386. THOMAS PE, WHARRAM BL, GOYAL M, WIGGINS JE, HOLZMAN LB, WIGGINS RC: GLEPP1, a renal glomerular epithelial cell (podocyte) membrane protein-tyrosine phosphatase: Identification, molecular cloning, and characterization in rabbit. *J Biol Chem* 269:19953–19962, 1994
  387. SHANKLAND SJ: Cell-cycle control and renal disease. *Kidney Int* 52:294–308, 1997
  388. SHERR CJ, ROBERTS JM: Inhibitors of mammalian G1 cyclin-dependent kinases. *Genes Dev* 9:1149–1163, 1995
  389. SHANKLAND SJ, PIPPIN J, FLANAGAN M, COATS SR, NANGAKU M, GORDON KL, ROBERTS JM, COUSER WG, JOHNSON RJ: Mesangial cell proliferation mediated by PDGF and bFGF is determined by levels of the cyclin kinase inhibitor p27<sup>Kip1</sup>. *Kidney Int* 51:1088–1099, 1997
  390. SHANKLAND SJ, HUGO C, COATS SR, NANGAKU M, PICHLER RH, GORDON KL, PIPPIN J, ROBERTS JM, COUSER WG, JOHNSON RJ: Changes in cell-cycle protein expression during experimental mesangial proliferative glomerulonephritis. *Kidney Int* 50:1230–1239, 1996
  391. OPHASCHAROENSUK V, FERO ML, HUGHES J, ROBERTS JM, SHANKLAND SJ: The cyclin dependent kinase inhibitor p27<sup>Kip1</sup> safeguards against inflammatory injury. *Nat Med* 4:575–580, 1998
  392. SHANKLAND SJ, FLOEGE J, THOMAS SE, NANGAKU M, HUGO C, PIPPIN J, HENNE K, HOCKENBERRY DM, JOHNSON RJ, COUSER WG: Cyclin kinase inhibitors are increased during experimental membranous nephropathy: Potential role in limiting glomerular epithelial cell proliferation in vivo. *Kidney Int* 52:404–413, 1997
  393. KAWAMURA T, YOSHIOKA T, BILLS T, FOGO A, ICHIKAWA I: Glucocorticoid activates glomerular antioxidant enzymes and protects glomeruli from oxidant injuries. *Kidney Int* 40:291–301, 1991
  394. KITAMURA M: Strategic gene transfer into the kidney: Current status and prospects. *Clin Exp Nephrol* 1:157–178, 1997
  395. KITAMURA M, FINE LG: Gene transfer into the kidney: Promises and limitations. *Kidney Int* 52:S86–S90, 1997
  396. TOMITA N, HIGAKI J, MORISHITA R, KATO K, MIKAMI H, KANEDA Y, OGIHARA T: Direct in vivo gene introduction into rat kidney. *Biochem Biophys Res Commun* 186:129–134, 1992
  397. IMAI E, ISAKA Y, AKAGI Y, KANEDA Y: Gene transfer into the glomerulus by hemagglutinating virus of Japan (HVJ)-liposome method, in *Genetic Manipulation of the Kidney*, edited by KITAMURA M. *Exp Nephrol* 5:112–117, 1997
  398. HEIKKILA P, PARPALA T, LUKKARINEN O, WEBER M, TRYGGVASON K: Adenovirus-mediated gene transfer into kidney glomeruli using an ex vivo and in vivo kidney perfusion system: First step towards gene therapy of Alport syndrome. *Gene Ther* 3:21–27, 1996
  399. TRYGGVASON K, HEIKKILA P, PETERSSON E, TIBELL A, THORNER P: Can Alport syndrome be treated by gene therapy? *Kidney Int* 51:1493–1499, 1997
  400. YOKOO T, KITAMURA M: Gene transfer of interleukin-1 receptor antagonist into the renal glomerulus via a mesangial cell vector. *Biochem Biophys Res Commun* 226:883–888, 1996
  401. KITAMURA M: Creation of a reversible on/off system for site-specific in vivo control of exogenous gene activity in the renal glomerulus. *Proc Natl Acad Sci USA* 93:7387–7391, 1996
  402. KITAMURA M, KAWACHI H: Creation of an in vivo cytosensor using engineered mesangial cells: Automatic sensing of glomerular inflammation controls transgene activity. *J Clin Invest* 100:1394–1399, 1997
  403. KOOYMAN DL, BYRNE GW, MCCLELLAN S, NIELSEN D, TONE M, WALDMANN H, COFFMAN TM, MCCURRY KR, PLATT JL, LOGAN JS: In vivo transfer of GPI-linked complement restriction factors from erythrocytes to the endothelium. *Science* 269:89–92, 1995
  404. MCCURRY KR, KOOYMAN DL, ALVARADO CG, COTTERELL AH, MARTIN MJ, LOGAN JS, PLATT JL: Human complement regulatory proteins protect swine-to-primate cardiac xenografts from humoral injury. *Nat Med* 1:423–427, 1995
  405. ZHU G, NICOLSON AG, ZHENG XX, STROM TB, SUKHATME VP: Adenovirus-mediated  $\beta$ -galactosidase gene delivery to the liver leads to protein deposition in kidney glomeruli. *Kidney Int* 52:992–999, 1997
  406. KAWACHI H, IWANAGA T, TOYABE S, OITE T, SHIMIZU F: Mesangial sclerotic change with persistent proteinuria in rats after two consecutive injection of monoclonal antibody 1-22-3. *Clin Exp Immunol* 90:129–134, 1992
  407. KASINATH BS: Glomerular endothelial cell proteoglycans: Regulation by TGF- $\beta$ 1. *Arch Biochem Biophys* 305:370–377, 1993
  408. FLOEGE J, JOHNSON RJ, GORDON K, IIDA H, PRITZL P, YOSHIMURA A, CAMPBELL C, ALPERS CE, COUSER WG: Increased synthesis of extracellular matrix in mesangial proliferative nephritis. *Kidney Int* 40:477–488, 1991
  409. MORIYAMA T, XIA C, MIYAI A, AKAGI Y, ISAKA Y, KANEKO T, KAWADA N, UEDA N, YAMAUCHI A, HORIO M, ANDO A, IMAI E, HORI M: Involvement of MAP kinase phosphatase-1 in mesangial cell growth regulation. *Nephrology* 3:443–453, 1997
  410. SUN H, CHARLES CH, LAU LF, TONKS NK: MKP-1 (3CH134), an immediate early gene product, is a dual specificity phosphatase that dephosphorylates MAP kinase in vivo. *Cell* 75:487–493, 1993
  411. FRANKLIN CC, KRAFT AS: Conditional expression of the mitogen-activated protein kinase (MAPK) phosphatase MKP-1 preferentially inhibits p38 MAPK and stress-activated protein kinase in U937 cells. *J Biol Chem* 272:16917–16923, 1997
  412. ANDERSON S, BRENNER BM: Intraglomerular hypertension: Implications and drug treatment. *Annu Rev Med* 39:243–253, 1988
  413. HARRIS RC, KONDO S, YASUDA T: Mechanical stress-cell function relationships in renal cells. *Exp Nephrol* 5:263–270, 1997